



An existence result for singular Finsler double phase problems

Csaba Farkas^a, Patrick Winkert^{b,*}

^a *Sapientia Hungarian University of Transylvania, Department of Mathematics and Computer Science, Târgu Mureș, Romania*

^b *Technische Universität Berlin, Institut für Mathematik, Straße des 17. Juni 136, 10623 Berlin, Germany*

Received 7 November 2020; accepted 16 March 2021

Abstract

In this paper, we study a class of singular double phase problems defined on Minkowski spaces in terms of Finsler manifolds and with right-hand sides that allow a certain type of critical growth for such problems. Under very general assumptions and based on variational tools we prove the existence of at least one non-trivial weak solution for such a problem. This is the first work dealing with a Finsler double phase operator even in the nonsingular case.

© 2021 Elsevier Inc. All rights reserved.

MSC: 35J15; 35J62; 58B20; 58J60

Keywords: Anisotropic double phase operator; Critical type exponent; Existence results; Minkowski space; Singular problems

1. Introduction

Let $\Omega \subset \mathbb{R}^N$, $N \geq 2$, be a bounded domain with Lipschitz boundary $\partial\Omega$ and let $F: \mathbb{R}^N \rightarrow [0, \infty)$ be a convex function of class $C^2(\mathbb{R}^N \setminus \{0\})$ which is absolutely homogeneous of degree one, that is,

* Corresponding author.

E-mail addresses: farkascs@ms.sapientia.ro, farkas.csaba2008@gmail.com (C. Farkas), winkert@math.tu-berlin.de (P. Winkert).

$$F(tx) = |t|F(x) \quad \text{for all } t \in \mathbb{R} \text{ and for all } x \in \mathbb{R}^N.$$

Then, one can consider the functional $E_F: H_0^1(\Omega) \rightarrow \mathbb{R}$ defined by

$$E_F(u) = \int_{\Omega} F^2(\nabla u) \, dx \quad \text{for } u \in H_0^1(\Omega),$$

which is the energy functional associated to the so-called Finsler Laplace operator given by

$$\Delta_F u = \operatorname{div}(F(\nabla u)\nabla F(\nabla u)).$$

This class of problems arises in mathematical physics where the objective is the minimization of $E_F: H_0^1(\Omega) \rightarrow \mathbb{R}$ under certain constraints (on perimeter or volume) while the solution corresponds to an optimal shape/configuration of anisotropic tension-surface. For instance, the minimization of the functional $E_F: H_0^1(\Omega) \rightarrow \mathbb{R}$ explains the specific polyhedral shape of crystal structures in solid crystals with sufficiently small grains, see Dinghas [17] and Taylor [45].

In the past years, many authors studied elliptic equations involving the Finsler Laplace operator. We refer, for instance, to the works of Cianchi-Salani [10] and Wang-Xia [46] who treated the Serrin-type overdetermined anisotropic problem involving the function F and the operator Δ_F while Farkas-Fodor-Kristály [21] considered a sublinear elliptic problem with Dirichlet boundary condition. Further studies of anisotropic phenomena can be found in the papers of Bellettini-Paolini [7], Belloni-Ferone-Kawohl [8], Della Pietra-Gavitone [16], Della Pietra-di Blasio-Gavitone [14], Della Pietra-Gavitone-Piscitelli [15] and Ferone-Kawohl [23]; see also the references therein. Nonlinear equations on general Finsler manifolds can be found in Farkas-Kristály-Varga [22], Farkas [20] and the references therein.

In this paper we are concerned with the following singular Finsler double phase problem given in the form

$$\begin{aligned} -\operatorname{div}\left(F^{p-1}(\nabla u)\nabla F(\nabla u) + \mu(x)F^{q-1}(\nabla u)\nabla F(\nabla u)\right) &= f(u) && \text{in } \Omega, \\ u &> 0 && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega, \end{aligned} \tag{1.1}$$

with

$$f(u) = u^{p^*-1} + \lambda\left(u^{\gamma-1} + g(u)\right),$$

where $\lambda > 0$ is a positive parameter and the following assumptions are supposed:

(H) (i) $0 < \gamma < 1$ and

$$2 \leq p < q < N, \quad \frac{q}{p} < 1 + \frac{1}{N}, \quad 0 \leq \mu(\cdot) \in C^{0,1}(\overline{\Omega}); \tag{1.2}$$

(ii) the function $g: \mathbb{R} \rightarrow \mathbb{R}$ is continuous and there exist $1 < \theta < p \leq \nu < p^*$ as well as nonnegative constants $c_i, i = 1, 2$ such that

$$g(s) \leq c_1 s^{\nu-1} + c_2 s^{\theta-1} \quad \text{for all } s \geq 0,$$

where p^* is the critical exponent to p given by

$$p^* := \frac{Np}{N-p}; \tag{1.3}$$

(iii) the function $F: \mathbb{R}^N \rightarrow [0, \infty)$ is a positively homogeneous Minkowski norm.

The differential operator in (1.1) is called the Finsler double phase operator. It is defined by

$$\operatorname{div} \left(F^{p-1}(\nabla u) \nabla F(\nabla u) + \mu(x) F^{q-1}(\nabla u) \nabla F(\nabla u) \right) \quad \text{for } u \in W_0^{1,\mathcal{H}}(\Omega), \tag{1.4}$$

where $W_0^{1,\mathcal{H}}(\Omega)$ is a suitable Musielak-Orlicz Sobolev space which is defined in Section 2. If $\mu \equiv 0$, the operator in (1.4) becomes the Finsler p -Laplacian (which reduces to the Finsler Laplacian in case $p = 2$) and if $\inf_{\Omega} \mu > 0$, we have the weighted Finsler (q, p) -Laplacian. Clearly, when $\mu \equiv 0$, (1.1) is strongly related to following Brezis-Nirenberg type problem

$$\begin{aligned} -\Delta_p u &= |u|^{p^*-2} u + \lambda g(u) && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega. \end{aligned}$$

Note that the Finsler p -Laplacian is often called anisotropic p -Laplacian. Since the word “anisotropic” also stands in the literature for problems where the exponent p is a function, we avoid this denotation and use only “Finsler”.

The operator (1.4) is related to the energy functional

$$\omega \mapsto \int_{\Omega} \left(F^p(\nabla \omega) + \mu(x) F^q(\nabla \omega) \right) dx, \tag{1.5}$$

where the integrand $H(x, \xi) = F^p(\xi) + \mu(x) F^q(\xi)$ for all $(x, \xi) \in \Omega \times \mathbb{R}^N$ has unbalanced growth, that is,

$$b_1 |\xi|^p \leq F^p(\xi) \leq H(x, \xi) \leq b_2 (1 + F^q(\xi)) \leq b_3 (1 + |\xi|^q)$$

for a. a. $x \in \Omega$ and for all $\xi \in \mathbb{R}^N$ with $b_1, b_2, b_3 > 0$. The integral functional (1.5) is characterized by the fact that the energy density changes its ellipticity and growth properties according to the point in the domain. To be more precise, the growth rate of (1.5) on the set $\{x \in \Omega : \mu(x) = 0\}$ is of order p and on the set $\{x \in \Omega : \mu(x) \neq 0\}$ it is of order q . So the integrand H switches between two different phases of elliptic behaviours.

When F coincide with the Euclidean norm, Zhikov [49] was the first who studied functionals of type (1.5) for models which describe strongly anisotropic materials, see also the monograph of Zhikov-Kozlov-Oleinik [50]. In the last decades, several authors investigated functionals of the form in (1.5) concerning regularity of (local) minimizers. We mention the papers of Baroni-Colombo-Mingione [3–5], Baroni-Kuusi-Mingione [6], Colombo-Mingione [12,13], Marcellini [32,31], Ok [35,36], Ragusa-Tachikawa [42] and the references therein.

Before we state the main result of the present paper, we recall the notion of the weak solution. A function $u \in W_0^{1,\mathcal{H}}(\Omega)$ is called a weak solution of problem (1.1) if $u^{\gamma-1}\varphi \in L^1(\Omega)$, $u > 0$ for a. a. $x \in \Omega$ and if

$$\begin{aligned} & \int_{\Omega} \left(F^{p-1}(\nabla u) \nabla F(\nabla u) + \mu(x) F^{q-1}(\nabla u) \nabla F(\nabla u) \right) \cdot \nabla \varphi \, dx \\ &= \int_{\Omega} u^{p^*-1} \varphi \, dx + \lambda \int_{\Omega} u^{\gamma-1} \varphi \, dx + \lambda \int_{\Omega} g(u) \varphi \, dx \end{aligned}$$

is satisfied for all $\varphi \in W_0^{1,\mathcal{H}}(\Omega)$.

Now, we are in the position to formulate our main result.

Theorem 1.1. *Let hypotheses (H) be satisfied. Then there exists $\lambda_* > 0$ such that for every $\lambda \in (0, \lambda_*)$, problem (1.1) has a nontrivial weak solution.*

Remark 1.2. Since we are looking for positive solutions and hypothesis (H)(ii) concerns the positive semiaxis $\mathbb{R}_+ = [0, +\infty)$, without any loss of generality, we may assume that $g(s) = 0$ for all $s \leq 0$.

To the best of our knowledge, this is the first paper which considers a double phase operator on a Minkowski space in the sense of Finsler geometry, see Bao–Chern–Shen [2]. Even if F is the Euclidean norm, there is only one work for singular double phase problems done by Chen–Ge–Wen–Cao [9] who proved the existence of a nontrivial weak solution with negative energy. Recent existence results for double phase problems in the case if F is the Euclidean norm and without singular term can be found, for example, in the papers of Colasuonno–Squassina [11], Liu–Dai [30], Gasiński–Papageorgiou [24], Gasiński–Winkert [25–27], Perera–Squassina [41] and the references therein.

So far, it is not clear what is the optimal critical exponent for Musielak–Orlicz Sobolev spaces even when F is the Euclidean norm, see, for example, the paper of Fan [18]. For the moment p^* seems to be best exponent (it is probably not optimal) and only a continuous (in general noncompact) embedding from $W_0^{1,\mathcal{H}}(\Omega) \hookrightarrow L^{p^*}(\Omega)$ is available. So we call it “type of critical growth”.

Finally, we mention recent papers which are very close to our topic. We refer to the works of Bahrouni–Rădulescu–Winkert [1], Marino–Winkert [33], Papageorgiou–Rădulescu–Repovš [37], Papageorgiou–Winkert [39,40], Zeng–Bai–Gasiński–Winkert [48,47] and the references therein.

The paper is organized as follows. In Section 2 we recall some basic properties of the Minkowski norm and introduce appropriate Musielak–Orlicz Sobolev spaces equipped with the Minkowski norm F . In Section 3 we first present the Minkowskian version of the inequality

$$|b|^r \geq |a|^r + r|a|^{r-2}a \cdot (b - a) + 2^{1-r}|a - b|^r \quad \text{for } r \geq 2 \text{ and for all } a, b \in \mathbb{R}^N,$$

see Lindqvist [29], which is the basis for the proof of the sequentially weakly lower semicontinuity of the corresponding energy functional to problem (1.1) on closed convex subsets of $W_0^{1,\mathcal{H}}(\Omega)$ not necessarily being balls. The sequentially weakly lower semicontinuity is one of

the main steps in order to prove our main result. At the end, based on the direct method of calculus along with variational tools, we are going to prove Theorem 1.1. Our approach follows ideas of the paper of Faraci-Farkas [19] in which the authors consider a singular problem driven by the p -Laplacian, that is, $\mu \equiv 0$ and $F(\xi) = \left(\sum_{i=1}^N |\xi_i|^2\right)^{1/2}$.

2. Preliminaries

In this section we will recall the main properties of the Minkowski space (\mathbb{R}^N, F) and the functional setting for the double phase operator.

Let $F: \mathbb{R}^N \rightarrow [0, \infty)$ be a positively homogeneous Minkowski norm, that is, F is a positive homogeneous function and verifies the properties

- $F \in C^\infty(\mathbb{R}^N \setminus \{0\})$;
- The Hessian matrix $\nabla^2(F^2/2)(x)$ is positive definite for all $x \neq 0$.

Note that the pair (\mathbb{R}^N, F) is the simplest not necessarily reversible Finsler manifold whose flag curvature is identically zero, the geodesics are straight lines and the intrinsic distance between two points $x, y \in \mathbb{R}^N$ is given by

$$d_F(x, y) = F(y - x).$$

In fact, (\mathbb{R}^N, d_F) is a quasi-metric space and in general, we have $d_F(x, y) \neq d_F(y, x)$.

A typical example for a Minkowski norm is the so called Randers metric which is defined by

$$F(x) = \sqrt{\langle Ax, x \rangle} + \langle b, x \rangle,$$

where A is a positive definite and symmetric $(N \times N)$ -type matrix and $b = (b_i) \in \mathbb{R}^N$ is a fixed vector such that $\sqrt{\langle A^{-1}b, b \rangle} < 1$. The pair (\mathbb{R}^N, F) is often called Randers space which describes the electromagnetic field of the physical space-time in general relativity, see Randers [43]. They are deduced as the solution of the Zermelo navigation problem.

In what follows we recall some basic properties of F , see Bao-Chern-Shen [2, §1.2].

Proposition 2.1. *Let $F: \mathbb{R}^N \rightarrow [0, \infty)$ be a positively homogeneous Minkowski norm. Then, the following assertions hold true:*

- (i) *Positivity: $F(x) > 0$ for all $x \neq 0$;*
- (ii) *Convexity: F and F^2 are strictly convex;*
- (iii) *Euler’s theorem: $x \cdot \nabla F(x) = F(x)$ and*

$$\nabla^2(F^2/2)(x)x \cdot x = F^2(x) \quad \text{for all } x \in \mathbb{R}^N \setminus \{0\};$$

- (iv) *Homogeneity: $\nabla F(tx) = \nabla F(x)$ and*

$$\nabla^2 F^2(tx) = \nabla^2 F^2(x) \quad \text{for all } x \in \mathbb{R}^N \setminus \{0\} \text{ and for all } t > 0.$$

For $1 \leq r < \infty$, we denote by $L^r(\Omega)$ and $L^r(\Omega; \mathbb{R}^N)$ the usual Lebesgue spaces equipped with the norm $\|\cdot\|_r$. If $1 < r < \infty$, then $W^{1,r}(\Omega)$ and $W_0^{1,r}(\Omega)$ stand for the Sobolev spaces endowed with the norms

$$\|u\|_{1,r,F} = \|F(\nabla u)\|_r + \|u\|_r,$$

and

$$\|u\|_{1,r,0,F} = \|F(\nabla u)\|_r,$$

respectively. Now, let $\mathcal{H}: \Omega \times [0, \infty) \rightarrow [0, \infty)$ be the function defined by

$$(x, t) \mapsto t^p + \mu(x)t^q,$$

where (1.2) is satisfied. The Musielak-Orlicz space $L^{\mathcal{H}}(\Omega)$ is defined by

$$L^{\mathcal{H}}(\Omega) = \left\{ u \mid u : \Omega \rightarrow \mathbb{R} \text{ is measurable and } \int_{\Omega} \mathcal{H}(x, |u|) \, dx < +\infty \right\}$$

equipped with the Luxemburg norm

$$\|u\|_{\mathcal{H}} = \inf \left\{ \tau > 0 : \rho_{\mathcal{H}}\left(\frac{u}{\tau}\right) \leq 1 \right\},$$

where

$$\rho_{\mathcal{H}}(u) := \int_{\Omega} \mathcal{H}(x, |u|) \, dx = \int_{\Omega} (|u|^p + \mu(x)|u|^q) \, dx. \tag{2.1}$$

We know that $L^{\mathcal{H}}(\Omega)$ is a reflexive Banach space.

The norm $\|\cdot\|_{\mathcal{H}}$ and the modular function $\rho_{\mathcal{H}}$ are related as follows, see Liu-Dai [30, Proposition 2.1].

Proposition 2.2. *Let (1.2) be satisfied, let $y \in L^{\mathcal{H}}(\Omega)$ and let $\rho_{\mathcal{H}}$ be defined by (2.1).*

- (i) *If $y \neq 0$, then $\|y\|_{\mathcal{H}} = \lambda$ if and only if $\rho_{\mathcal{H}}(\frac{y}{\lambda}) = 1$;*
- (ii) *$\|y\|_{\mathcal{H}} < 1$ (resp. $> 1, = 1$) if and only if $\rho_{\mathcal{H}}(y) < 1$ (resp. $> 1, = 1$);*
- (iii) *If $\|y\|_{\mathcal{H}} < 1$, then $\|y\|_{\mathcal{H}}^q \leq \rho_{\mathcal{H}}(y) \leq \|y\|_{\mathcal{H}}^p$;*
- (iv) *If $\|y\|_{\mathcal{H}} > 1$, then $\|y\|_{\mathcal{H}}^p \leq \rho_{\mathcal{H}}(y) \leq \|y\|_{\mathcal{H}}^q$;*
- (v) *$\|y\|_{\mathcal{H}} \rightarrow 0$ if and only if $\rho_{\mathcal{H}}(y) \rightarrow 0$;*
- (vi) *$\|y\|_{\mathcal{H}} \rightarrow +\infty$ if and only if $\rho_{\mathcal{H}}(y) \rightarrow +\infty$.*

The corresponding Sobolev space $W^{1,\mathcal{H}}(\Omega)$ is defined by

$$W^{1,\mathcal{H}}(\Omega) = \left\{ u \in L^{\mathcal{H}}(\Omega) : F(\nabla u) \in L^{\mathcal{H}}(\Omega) \right\}$$

equipped with the norm

$$\|u\|_{1,\mathcal{H},F} = \|F(\nabla u)\|_{\mathcal{H}} + \|u\|_{\mathcal{H}}.$$

By $W_0^{1,\mathcal{H}}(\Omega)$ we denote the completion of $C_0^\infty(\Omega)$ in $W^{1,\mathcal{H}}(\Omega)$ and thanks to (1.2) we have an equivalent norm on $W_0^{1,\mathcal{H}}(\Omega)$ given by

$$\|u\|_{1,\mathcal{H},0,F} = \|F(\nabla u)\|_{\mathcal{H}},$$

see Colasuonno-Squassina [11, Proposition 2.18]. Both spaces $W^{1,\mathcal{H}}(\Omega)$ and $W_0^{1,\mathcal{H}}(\Omega)$ are uniformly convex and so, reflexive Banach spaces.

Let $\rho_{\mathcal{H},F} : W_0^{1,\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$ be the corresponding modular to $W_0^{1,\mathcal{H}}(\Omega)$ defined by

$$\rho_{\mathcal{H},F}(u) := \int_{\Omega} (F^p(\nabla u) + \mu(x)F^q(\nabla u)) \, dx. \tag{2.2}$$

Similar to Proposition 2.2, we have the same relations between the norm $\|\cdot\|_{1,\mathcal{H},0,F}$ and the modular $\rho_{\mathcal{H},F}$.

Proposition 2.3. *Let (1.2) and (H)(iii) be satisfied and let $\rho_{\mathcal{H},F}$ be defined by (2.2). For $y \in W_0^{1,\mathcal{H}}(\Omega)$ we have the following assertions.*

- (i) *If $y \neq 0$, then $\|y\|_{1,\mathcal{H},0,F} = \lambda$ if and only if $\rho_{\mathcal{H},F}(\frac{y}{\lambda}) = 1$;*
- (ii) *$\|y\|_{1,\mathcal{H},0,F} < 1$ (resp. $> 1, = 1$) if and only if $\rho_{\mathcal{H},F}(y) < 1$ (resp. $> 1, = 1$);*
- (iii) *If $\|y\|_{1,\mathcal{H},0,F} < 1$, then $\|y\|_{1,\mathcal{H},0,F}^q \leq \rho_{\mathcal{H},F}(y) \leq \|y\|_{1,\mathcal{H},0,F}^p$;*
- (iv) *If $\|y\|_{1,\mathcal{H},0,F} > 1$, then $\|y\|_{1,\mathcal{H},0,F}^p \leq \rho_{\mathcal{H},F}(y) \leq \|y\|_{1,\mathcal{H},0,F}^q$;*
- (v) *$\|y\|_{1,\mathcal{H},0,F} \rightarrow 0$ if and only if $\rho_{\mathcal{H},F}(y) \rightarrow 0$;*
- (vi) *$\|y\|_{1,\mathcal{H},0,F} \rightarrow +\infty$ if and only if $\rho_{\mathcal{H},F}(y) \rightarrow +\infty$.*

In addition it is known that the embedding

$$W_0^{1,\mathcal{H}}(\Omega) \hookrightarrow L^r(\Omega)$$

is continuous whenever $r \leq p^*$ and compact whenever $r < p^*$, see Colasuonno-Squassina [11, Proposition 2.15]. Recall that p^* is the critical exponent to p , see (1.3).

Let $A : W_0^{1,\mathcal{H}}(\Omega) \rightarrow W_0^{1,\mathcal{H}}(\Omega)^*$ be the Finsler double phase operator defined by

$$\langle A(u), \varphi \rangle_{\mathcal{H},F} := \int_{\Omega} (F^{p-1}(\nabla u) \nabla F(\nabla u) + \mu(x)F^{q-1}(\nabla u) \nabla F(\nabla u)) \cdot \nabla \varphi \, dx \tag{2.3}$$

for all $u, \varphi \in W_0^{1,\mathcal{H}}(\Omega)$, where $\langle \cdot, \cdot \rangle_{\mathcal{H},F}$ is the duality pairing between $W_0^{1,\mathcal{H}}(\Omega)$ and its dual space $W_0^{1,\mathcal{H}}(\Omega)^*$. The properties of the operator $A : W_0^{1,\mathcal{H}}(\Omega) \rightarrow W_0^{1,\mathcal{H}}(\Omega)^*$ are summarized in the following proposition, see Liu-Dai [30], by taking the properties of F into account.

Proposition 2.4. *The operator A defined by (2.3) is bounded, continuous, monotone (hence maximal monotone) and of type (S_+) .*

3. Proof of the main result

In this section we present the proof of Theorem 1.1. We start with a lemma which is the Minkowskian version of the Lindqvist inequality; see Lindqvist [29]. Note that this inequality was proved for general Finsler manifolds by the first author [20]. For the sake of completeness we will also provide the proof here.

Lemma 3.1. *Let $F: \mathbb{R}^N \rightarrow [0, \infty)$ be a positively homogeneous Minkowski norm. For every $\xi, \beta \in \mathbb{R}^N$ and for every $r \geq 2$ we have*

$$F^r(\beta) \geq F^r(\xi) + rF^{r-1}(\xi)\nabla F(\xi) \cdot (\beta - \xi) + \frac{l_F^{\frac{r}{2}}}{2^{r-1}}F^r(\beta - \xi), \tag{3.1}$$

where

$$l_F = \min \left\{ \nabla^2 \left(\frac{F^2}{2} \right) (x)y \cdot y : F(x) = F(y) = 1 \right\}.$$

Proof. From the definition of l_F , one has that

$$F^2(t\xi + (1-t)\beta) \leq tF^2(\xi) + (1-t)F^2(\beta) - l_F t(1-t)F^2(\beta - \xi) \tag{3.2}$$

for all $\xi, \beta \in \mathbb{R}^N$ and for all $t \in [0, 1]$, see Farkas-Kristály-Varga [22, (2.19) on p. 1228] or Ohta [34]. Choosing $t = \frac{1}{2}$ in (3.2) gives

$$F^2 \left(\frac{\xi + \beta}{2} \right) \leq \frac{F^2(\xi) + F^2(\beta)}{2} - \frac{l_F}{4}F^2(\beta - \xi). \tag{3.3}$$

Note that for nonnegative $a, b \in \mathbb{R}$ we have

$$(a^r + b^r)^{\frac{1}{r}} = \left((a^r + b^r)^{\frac{2}{r}} \right)^{\frac{1}{2}} \leq \left(2^{\frac{2}{r}-1} (a^{r\frac{2}{r}} + b^{r\frac{2}{r}}) \right)^{\frac{1}{2}} \leq (a^2 + b^2)^{\frac{1}{2}}.$$

Applying the inequality above in combination with (3.3) leads to

$$\begin{aligned} F^r \left(\frac{\xi + \beta}{2} \right) + \frac{l_F^{\frac{r}{2}}}{2^r}F^r(\beta - \xi) &\leq \left(F^2 \left(\frac{\xi + \beta}{2} \right) + \frac{l_F}{4}F^2(\beta - \xi) \right)^{\frac{r}{2}} \\ &\leq \left(\frac{F^2(\xi) + F^2(\beta)}{2} \right)^{\frac{r}{2}}. \end{aligned} \tag{3.4}$$

By the convexity of the function $(\cdot)^{\frac{r}{2}}$ we have

$$\left(\frac{F^2(\xi) + F^2(\beta)}{2}\right)^{\frac{r}{2}} \leq \frac{1}{2}F^r(\xi) + \frac{1}{2}F^r(\beta). \tag{3.5}$$

On the other hand, by convexity, one has

$$F^r\left(\frac{\xi + \beta}{2}\right) \geq F^r(\xi) + \frac{r}{2}F^{r-1}(\xi)\nabla F(\xi) \cdot (\beta - \xi). \tag{3.6}$$

Combining (3.4), (3.5) and (3.6) yields

$$\frac{r}{2}F^{r-1}(\xi)\nabla F(\xi) \cdot (\beta - \xi) + \frac{l_F^{\frac{r}{2}}}{2^r}F^r(\beta - \xi) + \frac{1}{2}F^r(\xi) \leq \frac{1}{2}F^r(\beta),$$

for all $\xi, \beta \in \mathbb{R}^N$. Multiplying the last inequality by the factor 2 proves the assertion of the lemma. \square

Note that the constant l_F is the uniformity constant of F and $l_F \in [0, 1]$, see Farkas-Kristály-Varga [22] and Otha [34].

Let $J_\lambda : W_0^{1,\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$ be the energy functional corresponding to problem (1.1) given by

$$\begin{aligned} J_\lambda(u) = & \int_{\Omega} \left(\frac{1}{p}F^p(\nabla u) + \frac{\mu(x)}{q}F^q(\nabla u) \right) dx - \frac{1}{p^*} \int_{\Omega} (u_+)^{p^*} dx \\ & - \frac{\lambda}{\gamma} \int_{\Omega} (u_+)^{\gamma} dx - \lambda \int_{\Omega} G(u_+) dx, \end{aligned}$$

where $u_{\pm} = \max\{\pm u, 0\}$ and $G(s) = \int_0^s g(t) dt$. It is clear that J_λ does not belong to $C^1(W_0^{1,\mathcal{H}}(\Omega))$ due to the appearance of the singular term.

Furthermore we denote by κ_{p^*} the inverse of the Sobolev embedding constant of $W_0^{1,p}(\Omega) \hookrightarrow L^{p^*}(\Omega)$, that is,

$$(\kappa_{p^*})^{-1} = \inf_{\substack{u \in W_0^{1,p}(\Omega), \\ u \neq 0}} \frac{\|u\|_{1,p,0,F}}{\|u\|_{p^*}}.$$

Now, we are going to prove that the energy functional $J_\lambda : W_0^{1,\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$ is sequentially weakly lower semicontinuous on closed convex subsets of $W_0^{1,\mathcal{H}}(\Omega)$.

Proposition 3.2. *For every $\lambda > 0$ and for every $\sigma \in (0, \sigma^*)$, where*

$$\sigma^* := \left(\frac{p^*}{p} \frac{l_F^{\frac{p}{2}}}{2^{p-1}\kappa_{p^*}^{p^*}} \right)^{\frac{1}{p^*-p}},$$

the restriction of J_λ to the closed convex set B_σ which is given by

$$B_\sigma := \left\{ u \in W_0^{1,\mathcal{H}}(\Omega) : \|u\|_{1,p,0,F} \leq \sigma \right\},$$

is sequentially weakly lower semicontinuous.

Proof. Let $\lambda > 0$ and let $\sigma \in (0, \sigma^*)$. We are going to prove that the functional $J_\lambda: W_0^{1,\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$ is weakly lower semicontinuous on B_σ . To this end, let $\{u_n\}_{n \in \mathbb{N}} \subset B_\sigma$ be such that $u_n \rightharpoonup u$ in $W_0^{1,\mathcal{H}}(\Omega)$ and let

$$I(u) = \int_\Omega \left(\frac{1}{p} F^p(\nabla u) + \frac{\mu(x)}{q} F^q(\nabla u) \right) dx - \frac{1}{p^*} \int_\Omega (u_+)^{p^*} dx.$$

Note that, up to a subsequence if necessary, we have for $r < p^*$, $u_n \rightarrow u$ in $L^r(\Omega)$ and $\nabla u_n \rightharpoonup \nabla u$ in $L^p(\Omega)$. Therefore, it suffices to prove that

$$\liminf_{n \rightarrow \infty} (I(u_n) - I(u)) \geq 0,$$

see hypothesis (H)(ii).

Applying (3.1) we get the following inequalities

$$\begin{aligned} & \int_\Omega \frac{1}{p} F^p(\nabla u_n) dx \\ & \geq \int_\Omega \frac{1}{p} F^p(\nabla u) dx + \int_\Omega F^{p-1}(\nabla u) \nabla F(\nabla u) \cdot \nabla (u_n - u) dx + \frac{l_F^{\frac{p}{2}}}{p2^{p-1}} \int_\Omega F^p(\nabla u_n - \nabla u) dx, \end{aligned}$$

and

$$\begin{aligned} \int_\Omega \frac{\mu(x)}{q} F^q(\nabla u_n) dx & \geq \int_\Omega \frac{\mu(x)}{q} F^q(\nabla u) dx + \int_\Omega \mu(x) F^{q-1}(\nabla u) \nabla F(\nabla u) \cdot \nabla (u_n - u) dx \\ & \quad + \frac{l_F^{\frac{q}{2}}}{q2^{q-1}} \int_\Omega \mu(x) F^q(\nabla u_n - \nabla u) dx. \end{aligned}$$

On the other hand, by the Brezis-Lieb lemma, see, for example, Papageorgiou-Winkert [38, Lemma 4.1.22], we have

$$\liminf_{n \rightarrow \infty} \left(\int_\Omega |u_n|^{p^*} dx - \int_\Omega |u|^{p^*} dx \right) = \liminf_{n \rightarrow \infty} \int_\Omega |u_n - u|^{p^*} dx.$$

Therefore since

$$\begin{aligned} &\int_{\Omega} F^{p-1}(\nabla u)\nabla F(\nabla u) \cdot \nabla (u_n - u) \, dx \rightarrow 0, \\ &\int_{\Omega} \mu(x)F^{q-1}(\nabla u)\nabla F(\nabla u) \cdot \nabla (u_n - u) \, dx \rightarrow 0, \\ &\frac{l_F^{\frac{q}{2}}}{q2^{q-1}} \int_{\Omega} \mu(x)F^q(\nabla u_n - \nabla u) \, dx \geq 0, \end{aligned}$$

one has

$$\begin{aligned} \liminf_{n \rightarrow \infty} (I(u_n) - I(u)) &\geq \liminf_{n \rightarrow \infty} \left(\frac{l_F^{\frac{p}{2}}}{p2^{p-1}} \int_{\Omega} F^p(\nabla u_n - \nabla u) \, dx - \frac{1}{p^*} \int_{\Omega} |u_n - u|^{p^*} \, dx \right) \\ &\geq \liminf_{n \rightarrow \infty} \left(\frac{l_F^{\frac{p}{2}}}{p2^{p-1}} \|u_n - u\|_{1,p,0,F}^p - \frac{1}{p^*} k_{p^*}^{p^*} \|u_n - u\|_{1,p,0,F}^{p^*} \right) \\ &\geq \liminf_{n \rightarrow \infty} \|u_n - u\|_{1,p,0,F}^p \left(\frac{l_F^{\frac{p}{2}}}{p2^{p-1}} - \frac{1}{p^*} k_{p^*}^{p^*} \sigma^{p^*-p} \right) \geq 0, \end{aligned}$$

which proves the assertion of the proposition. \square

For further use, we introduce the functionals $I_1 : W_0^{1,\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$ and $I_2 : L^{\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$ defined by

$$I_1(u) = -\frac{1}{q} \int_{\Omega} \mu(x)F^q(\nabla u) \, dx + \frac{1}{p^*} \int_{\Omega} (u_+)^{p^*} \, dx + \frac{\lambda}{\gamma} \int_{\Omega} (u_+)^{\gamma} \, dx + \lambda \int_{\Omega} G(u_+) \, dx,$$

and

$$I_2(u) = \frac{1}{p^*} \int_{\Omega} (u_+)^{p^*} \, dx + \frac{\lambda}{\gamma} \int_{\Omega} (u_+)^{\gamma} \, dx + \lambda \int_{\Omega} G(u_+) \, dx.$$

Now we are in the position to prove Theorem 1.1.

Proof of Theorem 1.1. Let $\lambda > 0$ and let $\sigma \in (0, \sigma^*)$, where σ^* is as in Proposition 3.2. We define

$$\varphi_{\lambda}(\sigma) := \inf_{\|u\|_{1,p,0,F} < \sigma} \frac{\sup_{B_{\sigma}} I_1 - I_1(u)}{\sigma^p - \|u\|_{1,p,0,F}^p}$$

and

$$\psi_\lambda(\sigma) := \sup_{B_\sigma} I_1.$$

We claim that there exist $\lambda, \sigma > 0$ small enough such that

$$\varphi_\lambda(\sigma) < \frac{1}{p}. \tag{3.7}$$

We claim that (3.7) holds if there exist $\lambda, \sigma > 0$ such that

$$\inf_{\xi < \sigma} \frac{\psi_\lambda(\sigma) - \psi_\lambda(\xi)}{\sigma^p - \xi^p} < \frac{1}{p}. \tag{3.8}$$

Note that, by taking $\xi = \sigma - \varepsilon$ for some $\varepsilon \in (0, \sigma)$, it follows

$$\begin{aligned} \frac{\psi_\lambda(\sigma) - \psi_\lambda(\xi)}{\sigma^p - \xi^p} &= \frac{\psi_\lambda(\sigma) - \psi_\lambda(\sigma - \varepsilon)}{\sigma^p - (\sigma - \varepsilon)^p} \\ &= \frac{\psi_\lambda(\sigma) - \psi_\lambda(\sigma - \varepsilon)}{\varepsilon} \cdot \frac{-\frac{\varepsilon}{\sigma}}{\sigma^{p-1}[(1 - \frac{\varepsilon}{\sigma})^p - 1]}. \end{aligned}$$

So passing to the limit as $\varepsilon \rightarrow 0$, we get that (3.8) is fulfilled if

$$\limsup_{\varepsilon \rightarrow 0^+} \frac{\psi_\lambda(\sigma) - \psi_\lambda(\sigma - \varepsilon)}{\varepsilon} < \sigma^{p-1}. \tag{3.9}$$

First note that

$$\begin{aligned} \frac{1}{\varepsilon} |\psi_\lambda(\sigma) - \psi_\lambda(\sigma - \varepsilon)| &= \frac{1}{\varepsilon} \left| \sup_{v \in B_1} I_1(\sigma v) - \sup_{v \in B_1} I_1((\sigma - \varepsilon)v) \right| \\ &\leq \frac{1}{\varepsilon} \sup_{v \in B_1} |I_1(\sigma v) - I_1((\sigma - \varepsilon)v)| \\ &\leq \frac{1}{\varepsilon} \sup_{v \in B_1} \left| \frac{(\sigma - \varepsilon)^q - \sigma^q}{q} \int_{\Omega} \mu(x) F^q(\nabla v) \, dx + I_2(\sigma v) - I_2((\sigma - \varepsilon)v) \right|. \end{aligned}$$

Applying (H)(ii) and the continuous embedding $W_0^{1,p}(\Omega) \hookrightarrow L^{p^*}(\Omega)$ we obtain

$$\begin{aligned} \frac{\psi_\lambda(\sigma) - \psi_\lambda(\sigma - \varepsilon)}{\varepsilon} &\leq \frac{1}{\varepsilon} \sup_{\|v\|_{1,p,0,F} \leq 1} \int_{\Omega} \left| \int_{(\sigma - \varepsilon)v_+(x)}^{\sigma v_+(x)} \left[t^{p^*-1} + \lambda t^{\gamma-1} + \lambda g(t) \right] dt \right| dx \\ &\leq \frac{\kappa_{p^*}^{p^*}}{p^*} \left| \frac{\sigma^{p^*} - (\sigma - \varepsilon)^{p^*}}{\varepsilon} \right| + \lambda \frac{\kappa_{p^*}^\gamma |\Omega|^{\frac{p^*-\gamma}{p^*}}}{\gamma} \left| \frac{\sigma^\gamma - (\sigma - \varepsilon)^\gamma}{\varepsilon} \right| \end{aligned}$$

$$+ \lambda c_1 \frac{\kappa_{p^*}^\nu |\Omega|^{\frac{p^*-\nu}{p^*}}}{\nu} \left| \frac{\sigma^\nu - (\sigma - \varepsilon)^\nu}{\varepsilon} \right| + \lambda c_2 \frac{\kappa_{p^*}^\theta |\Omega|^{\frac{p^*-\theta}{p^*}}}{\theta} \left| \frac{\sigma^\theta - (\sigma - \varepsilon)^\theta}{\varepsilon} \right|.$$

Therefore

$$\begin{aligned} & \limsup_{\varepsilon \rightarrow 0^+} \frac{\psi_\lambda(\sigma) - \psi_\lambda(\sigma - \varepsilon)}{\varepsilon} \\ & \leq \kappa_{p^*}^{p^*} \sigma^{p^*-1} + \lambda \kappa_{p^*}^\gamma |\Omega|^{\frac{p^*-\gamma}{p^*}} \sigma^{\gamma-1} + \lambda c_1 \kappa_{p^*}^\nu |\Omega|^{\frac{p^*-\nu}{p^*}} \sigma^{\nu-1} + \lambda c_2 \kappa_{p^*}^\theta |\Omega|^{\frac{p^*-\theta}{p^*}} \sigma^{\theta-1}. \end{aligned}$$

We now consider the function $\Lambda : (0, +\infty) \rightarrow \mathbb{R}$ defined by

$$\Lambda(s) = \frac{s^{p-\gamma} - \kappa_{p^*}^{p^*} s^{p^*-\gamma}}{\kappa_{p^*}^\gamma |\Omega|^{\frac{p^*-\gamma}{p^*}} + c_1 \kappa_{p^*}^\nu |\Omega|^{\frac{p^*-\nu}{p^*}} s^{\nu-\gamma} + c_2 \kappa_{p^*}^\theta |\Omega|^{\frac{p^*-\theta}{p^*}} s^{\theta-\gamma}}.$$

Applying L'Hospital's rule, it is clear that $\lim_{s \rightarrow 0} \Lambda(s) = 0$ while $\lim_{s \rightarrow \infty} \Lambda(s) = -\infty$. On the other hand, there exists $s_0 > 0$ small enough such that $\Lambda(s) > 0$ for every $s \in (0, s_0)$. Thus, there exists $s_{\max} > 0$ such that

$$\Lambda(s_{\max}) = \max_{s>0} \Lambda(s).$$

Therefore we may choose

$$\lambda_* := \Lambda(\min\{s_{\max}, \sigma^*\}).$$

Hence, if $\lambda < \lambda_*$ and $\sigma < \min\{s_{\max}, \sigma^*\}$, we have that (3.9) and so (3.7) are fulfilled.

From (3.7) we know that we can find $u \in W_0^{1,\mathcal{H}}(\Omega)$ with $\|u\|_{1,p,0,F} \leq \sigma$ such that

$$J_\lambda(u) < \frac{1}{p} \sigma^p - I_1(u_1) \quad \text{for all } u_1 \in B_\sigma. \tag{3.10}$$

Since $J_\lambda|_{B_\sigma}$ is sequentially weakly lower semicontinuous by Proposition 3.2, the energy functional $J_\lambda : W_0^{1,\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$ restricted to B_σ has a global minimizer $u_* \in W_0^{1,\mathcal{H}}(\Omega)$ with $\|u_*\|_{1,p,0,F} \leq \sigma$. If $\|u_*\|_{1,p,0,F} = \sigma$, then from (3.10) we obtain

$$J_\lambda(u_*) = \frac{1}{p} \sigma^p - I_1(u_*) > J_\lambda(u),$$

which is a contradiction. It follows that u_* is a local minimum for J_λ with $\|u_*\|_{1,p,0,F} < \sigma$. In summary, the energy functional J_λ has a local minimum $u =: u_* \in B_\sigma$ for $\lambda < \lambda_*$.

Due to the presence of the singular term, zero is not a local minimizer of J_λ . Indeed, if $v \in W_0^{1,\mathcal{H}}(\Omega)$ is positive in Ω and $\tau > 0$, then,

$$\begin{aligned}
 J_\lambda(\tau v) &\leq \frac{\tau^p}{p} \|v\|_{1,p,0,F}^p + \frac{\tau^q}{q} \int_\Omega \mu(x) F^q(\nabla v) \, dx - \frac{\tau^{p^*}}{p^*} \|v\|_{p^*}^{p^*} - \lambda \frac{\tau^\gamma}{\gamma} \|v\|_\gamma^\gamma \\
 &\quad + \lambda \frac{c_1 \tau^\nu}{\nu} \|v\|_\nu^\nu + \lambda \frac{c_2 \tau^\theta}{\theta} \|v\|_\theta^\theta.
 \end{aligned}$$

Thus $J_\lambda(\tau v) < 0$ for sufficiently small $\tau > 0$.

It remains to prove that u is a weak solution of (1.1). First, we prove that $u > 0$ a. e. in Ω . To this end, for $t > 0$ small enough, one has $u + tu_- \in B_\sigma$ and $(u + tu_-)_+ = u_+$. Thus,

$$\begin{aligned}
 0 &\leq \frac{J_\lambda(u + tu_-) - J_\lambda(u)}{t} \\
 &= \frac{1}{p} \left(\frac{\|u + tu_- - \|u\|_{1,p,0,F}^p}{t} - \|u\|_{1,p,0,F}^p \right) + \frac{1}{q} \int_\Omega \mu(x) \frac{F^q(\nabla(u + tu_-)) - F^q(\nabla u)}{t} \, dx \\
 &\rightarrow \int_\Omega F^{p-1}(\nabla u) \nabla F(\nabla u) \cdot \nabla u_- \, dx + \int_\Omega \mu(x) F^{q-1}(\nabla u) \nabla F(\nabla u) \cdot \nabla u_- \, dx \quad \text{as } t \rightarrow 0^+.
 \end{aligned}$$

On the other hand, using Proposition 2.1 (iii), one has that

$$\int_\Omega F^{p-1}(\nabla u) \nabla F(\nabla u) \cdot \nabla u_- \, dx = - \int_\Omega F^{p-1}(\nabla u_-) \nabla F(\nabla u_-) \cdot \nabla u_- \, dx = - \int_\Omega F^p(\nabla u_-) \, dx$$

and

$$\int_\Omega \mu(x) F^{q-1}(\nabla u) \nabla F(\nabla u) \cdot \nabla u_- \, dx = - \int_\Omega \mu(x) F^q(\nabla u_-) \, dx.$$

Therefore

$$0 \leq \lim_{t \rightarrow 0} \frac{J_\lambda(u + tu_-) - J_\lambda(u)}{t} = - \int_\Omega F^p(\nabla u_-) \, dx - \int_\Omega \mu(x) F^q(\nabla u_-) \, dx,$$

which shows that $u_- = 0$ and so $u \geq 0$ a. e. in Ω .

Assume that there exists a set A of positive measure such that $u = 0$ in A . Let $\varphi: \Omega \rightarrow \mathbb{R}$ be a function in $W_0^{1,\mathcal{H}}(\Omega)$ which is positive in Ω . For $t > 0$ small enough, we know that $u + t\varphi \in B_\sigma$ and $(u + t\varphi)^\gamma > u^\gamma$ a. e. in Ω . Hence, we obtain

$$\begin{aligned}
 0 &\leq \frac{J_\lambda(u + t\varphi) - J_\lambda(u)}{t} \\
 &= \frac{1}{p} \left(\frac{\|u + t\varphi\|_{1,p,0,F}^p - \|u\|_{1,p,0,F}^p}{t} \right) + \frac{1}{q} \int_\Omega \mu(x) \frac{F^q(\nabla(u + t\varphi))^q - F^q(\nabla u)}{t} dx \\
 &\quad - \frac{1}{p^*} \int_\Omega \frac{(u + t\varphi)^{p^*} - u^{p^*}}{t} dx - \frac{\lambda}{\gamma t^{1-\gamma}} \int_A \varphi^\gamma dx \\
 &\quad - \frac{\lambda}{\gamma} \int_{\Omega \setminus A} \frac{(u + t\varphi)^\gamma - u^\gamma}{t} dx - \lambda \int_\Omega \frac{G((u + t\varphi)) - G(u)}{t} dx \\
 &< \frac{1}{p} \left(\frac{\|u + t\varphi\|_{1,p,0,F}^p - \|u\|_{1,p,0,F}^p}{t} \right) + \frac{1}{q} \int_\Omega \mu(x) \frac{F^q(\nabla(u + t\varphi))^q - F^q(\nabla u)}{t} dx \\
 &\quad - \frac{1}{p^*} \int_\Omega \frac{(u + t\varphi)^{p^*} - u^{p^*}}{t} dx - \frac{\lambda}{\gamma t^{1-\gamma}} \int_A \varphi^\gamma dx - \lambda \int_\Omega \frac{G((u + t\varphi)) - G(u)}{t} dx.
 \end{aligned}$$

This gives

$$0 \leq \frac{J_\lambda(u + t\varphi) - J_\lambda(u)}{t} \longrightarrow -\infty \text{ as } t \rightarrow 0^+, \tag{3.11}$$

which is a contradiction. Thus, $u > 0$ a. e. in Ω .

In the next step we are going to prove that

$$u^{\gamma-1}\varphi \in L^1(\Omega) \text{ for all } \varphi \in W_0^{1,\mathcal{H}}(\Omega) \tag{3.12}$$

and

$$\begin{aligned}
 &\int_\Omega \left(F(\nabla u)^{p-1} + \mu(x) F(\nabla u)^{q-1} \right) \nabla F(\nabla u) \cdot \nabla \varphi dx \\
 &\quad - \int_\Omega u^{p^*-1} \varphi dx - \lambda \int_\Omega u^{\gamma-1} \varphi dx - \lambda \int_\Omega g(u) \varphi dx \geq 0
 \end{aligned} \tag{3.13}$$

for all $\varphi \in W_0^{1,\mathcal{H}}(\Omega)$ with $\varphi \geq 0$. Choose $\varphi \in W_0^{1,\mathcal{H}}(\Omega)$ with $\varphi \geq 0$ and fix a decreasing sequence $\{t_n\}_{n \in \mathbb{N}} \subseteq (0, 1]$ such that $\lim_{n \rightarrow \infty} t_n = 0$. For $n \in \mathbb{N}$, the functions

$$h_n(x) = \frac{(u(x) + t_n\varphi(x))^\gamma - u(x)^\gamma}{t_n}$$

are measurable, nonnegative and we have

$$\lim_{n \rightarrow \infty} h_n(x) = \gamma u(x)^{\gamma-1} \varphi(x) \text{ for a. a. } x \in \Omega.$$

From Fatou’s lemma we infer

$$\int_{\Omega} u^{\gamma-1} \varphi \, dx \leq \frac{1}{\gamma} \liminf_{n \rightarrow \infty} \int_{\Omega} h_n \, dx. \tag{3.14}$$

Similar to (3.11), we have for $n \in \mathbb{N}$ large enough

$$\begin{aligned} 0 &\leq \frac{J_{\lambda}(u + t_n \varphi) - J_{\lambda}(u)}{t_n} \\ &= \frac{1}{p} \frac{\|u + t_n \varphi\|_{1,p,0,F}^p - \|u\|_{1,p,0,F}^p}{t_n} + \frac{1}{q} \int_{\Omega} \mu(x) \frac{F^q(\nabla(u + t\varphi))^q - F^q(\nabla u)}{t} \, dx \\ &\quad - \frac{1}{p^*} \int_{\Omega} \frac{(u + t_n \varphi)^{p^*} - u^{p^*}}{t_n} \, dx - \frac{\lambda}{\gamma} \int_{\Omega} h_n \, dx - \lambda \int_{\Omega} \frac{G(u + t_n \varphi) - G(u)}{t_n} \, dx. \end{aligned}$$

Passing to the limit as $n \rightarrow \infty$ and applying (3.14) we have (3.12) and we get that

$$\begin{aligned} \lambda \int_{\Omega} u^{\gamma-1} \varphi \, dx &\leq \int_{\Omega} \left(F(\nabla u)^{p-1} + \mu(x) F(\nabla u)^{q-1} \right) \nabla F(\nabla u) \cdot \nabla \varphi \, dx \\ &\quad - \int_{\Omega} u^{p^*-1} \varphi \, dx - \lambda \int_{\Omega} g(u) \varphi \, dx, \end{aligned}$$

which shows (3.13). Note that it is enough to show (3.12) for nonnegative $\varphi \in W_0^{1,\mathcal{H}}(\Omega)$.

Now, let $\varepsilon \in (0, 1)$ be such that $(1 + t)u \in B_{\sigma}$ for all $t \in [-\varepsilon, \varepsilon]$. Note that the function $\alpha(t) := J_{\lambda}((1 + t)u)$ has a local minimum in zero. Using again Proposition 2.1 (iii) gives

$$\begin{aligned} 0 = \alpha'(0) &= \lim_{t \rightarrow 0} \frac{J_{\lambda}((1 + t)u) - J_{\lambda}(u)}{t} \\ &= \rho_{\mathcal{H},F}(u) - \int_{\Omega} u^{p^*} \, dx - \lambda \int_{\Omega} u^{\gamma} \, dx - \lambda \int_{\Omega} g(u)u \, dx. \end{aligned} \tag{3.15}$$

It remains to show that u is indeed a positive weak solution of (1.1). To this end we use the method of Lair-Shaker, see, for example, Lair-Shaker [28] or Sun-Wu-Long [44]. Let $\varphi \in W_0^{1,\mathcal{H}}(\Omega)$ and take the test function $v = (u + \varepsilon\varphi)_+$ in (3.13). Applying (3.15) we have that

$$\begin{aligned} 0 &\leq \int_{\{u+\varepsilon\varphi \geq 0\}} \left(F^{p-1}(\nabla u) + \mu(x) F^{q-1}(\nabla u) \right) \nabla F(\nabla u) \cdot \nabla(u + \varepsilon\varphi) \, dx \\ &\quad - \int_{\Omega} u^{p^*-1}(u + \varepsilon\varphi) \, dx - \lambda \int_{\Omega} u^{\gamma-1}(u + \varepsilon\varphi) \, dx - \lambda \int_{\Omega} g(u)(u + \varepsilon\varphi) \, dx \\ &= \rho_{\mathcal{H},F}(u) + \varepsilon \int_{\Omega} F^{p-1}(\nabla u) \nabla F(\nabla u) \cdot \nabla \varphi \, dx + \varepsilon \int_{\Omega} \mu(x) F^{q-1}(\nabla u) \nabla F(\nabla u) \cdot \nabla \varphi \, dx \end{aligned}$$

$$\begin{aligned}
 & - \int_{\Omega} u^{p^*} \, dx - \varepsilon \int_{\Omega} u^{p^*-1} \varphi \, dx - \lambda \int_{\Omega} u^{\gamma} \, dx - \varepsilon \lambda \int_{\Omega} u^{\gamma-1} \, dx \\
 & - \lambda \int_{\Omega} g(u)u \, dx - \varepsilon \lambda \int_{\Omega} g(u)\varphi \, dx - \int_{\{u+\varepsilon\varphi<0\}} F^p(\nabla u) \, dx \\
 & - \varepsilon \int_{\{u+\varepsilon\varphi<0\}} F^{p-1}(\nabla u)\nabla F(\nabla u) \cdot \nabla \varphi \, dx - \int_{\{u+\varepsilon\varphi<0\}} \mu(x)F^q(\nabla u) \, dx \tag{3.16} \\
 & - \varepsilon \int_{\{u+\varepsilon\varphi<0\}} \mu(x)F^{q-1}(\nabla u)\nabla F(\nabla u) \cdot \nabla \varphi \, dx + \int_{\{u+\varepsilon\varphi<0\}} u^{p^*-1}(u + \varepsilon\varphi) \, dx \\
 & + \lambda \int_{\{u+\varepsilon\varphi<0\}} u^{\gamma-1}(u + \varepsilon\varphi) \, dx + \lambda \int_{\{u+\varepsilon\varphi<0\}} g(u)(u + \varepsilon\varphi) \, dx \\
 & \leq \varepsilon \left[\int_{\Omega} \left(F^{p-1}(\nabla u) + \mu(x)F^{q-1}(\nabla u) \right) \nabla F(\nabla u) \cdot \nabla \varphi \, dx - \int_{\Omega} u^{p^*-1} \varphi \, dx \right. \\
 & \quad \left. - \lambda \int_{\Omega} u^{\gamma-1} \varphi \, dx - \lambda \int_{\Omega} g(u)\varphi \, dx \right] - \varepsilon \int_{\{u+\varepsilon\varphi<0\}} F^{p-1}(\nabla u)\nabla F(\nabla u) \cdot \nabla \varphi \, dx \\
 & - \varepsilon \int_{\{u+\varepsilon\varphi<0\}} \mu(x)F^{q-1}(\nabla u)\nabla F(\nabla u) \cdot \nabla \varphi \, dx.
 \end{aligned}$$

Note that the measure of the set $\{u + \varepsilon\varphi < 0\}$ goes to 0 as $\varepsilon \rightarrow 0$. Hence,

$$\int_{\{u+\varepsilon\varphi<0\}} F^{p-1}(\nabla u)\nabla F(\nabla u)\nabla \varphi \, dx + \int_{\{u+\varepsilon\varphi<0\}} \mu(x)F^{q-1}(\nabla u)\nabla F(\nabla u)\nabla \varphi \, dx \rightarrow 0.$$

Using this, dividing (3.16) by ε and passing to the limit as $\varepsilon \rightarrow 0$ we conclude that

$$\begin{aligned}
 & \int_{\Omega} \left(F(\nabla u)^{p-1} + \mu(x)F(\nabla u)^{q-1} \right) \nabla F(\nabla u) \cdot \nabla \varphi \, dx \\
 & - \int_{\Omega} u^{p^*-1} \varphi \, dx - \lambda \int_{\Omega} u^{\gamma-1} \varphi \, dx - \lambda \int_{\Omega} g(u)\varphi \, dx \geq 0.
 \end{aligned}$$

As φ was arbitrary chosen, we get from the last inequality that $u \in W_0^{1,\mathcal{H}}(\Omega)$ is a weak solution of (1.1). This finishes the proof. \square

Acknowledgment

C. Farkas was supported by the National Research, Development and Innovation Fund of Hungary, financed under the K_18 funding scheme, Project No. 127926 and by the Sapientia Foundation - Institute for Scientific Research, Romania, Project No. 17/11.06.2019.

References

- [1] A. Bahrouni, V.D. Rădulescu, P. Winkert, Double phase problems with variable growth and convection for the Baouendi–Grushin operator, *Z. Angew. Math. Phys.* 71 (6) (2020) 183.
- [2] D. Bao, S.-S. Chern, Z. Shen, *An Introduction to Riemann–Finsler Geometry*, Springer-Verlag, New York, 2000.
- [3] P. Baroni, M. Colombo, G. Mingione, Harnack inequalities for double phase functionals, *Nonlinear Anal.* 121 (2015) 206–222.
- [4] P. Baroni, M. Colombo, G. Mingione, Non-autonomous functionals, borderline cases and related function classes, *St. Petersburg Math. J.* 27 (2016) 347–379.
- [5] P. Baroni, M. Colombo, G. Mingione, Regularity for general functionals with double phase, *Calc. Var. Partial Differ. Equ.* 57 (2) (2018) 62.
- [6] P. Baroni, T. Kuusi, G. Mingione, Borderline gradient continuity of minima, *J. Fixed Point Theory Appl.* 15 (2) (2014) 537–575.
- [7] G. Bellettini, M. Paolini, Anisotropic motion by mean curvature in the context of Finsler geometry, *Hokkaido Math. J.* 25 (3) (1996) 537–566.
- [8] M. Belloni, V. Ferone, B. Kawohl, Isoperimetric inequalities, Wulff shape and related questions for strongly non-linear elliptic operators, *Z. Angew. Math. Phys.* 54 (5) (2003) 771–783.
- [9] Z.-Y. Chen, B. Ge, W.-S. Yuan, X.-F. Cao, Existence of solution for double-phase problem with singular weights, *Adv. Math. Phys.* 2020 (2020) 5376013.
- [10] A. Cianchi, P. Salani, Overdetermined anisotropic elliptic problems, *Math. Ann.* 345 (4) (2009) 859–881.
- [11] F. Colasuonno, M. Squassina, Eigenvalues for double phase variational integrals, *Ann. Mat. Pura Appl.* (4) 195 (6) (2016) 1917–1959.
- [12] M. Colombo, G. Mingione, Bounded minimisers of double phase variational integrals, *Arch. Ration. Mech. Anal.* 218 (1) (2015) 219–273.
- [13] M. Colombo, G. Mingione, Regularity for double phase variational problems, *Arch. Ration. Mech. Anal.* 215 (2) (2015) 443–496.
- [14] F. Della Pietra, G. di Blasio, N. Gavitone, Sharp estimates on the first Dirichlet eigenvalue of nonlinear elliptic operators via maximum principle, *Adv. Nonlinear Anal.* 9 (1) (2020) 278–291.
- [15] F. Della Pietra, N. Gavitone, G. Piscitelli, On the second Dirichlet eigenvalue of some nonlinear anisotropic elliptic operators, *Bull. Sci. Math.* 155 (2019) 10–32.
- [16] F. Della Pietra, N. Gavitone, Sharp estimates and existence for anisotropic elliptic problems with general growth in the gradient, *Z. Anal. Anwend.* 35 (1) (2016) 61–80.
- [17] A. Dinghas, Über einen geometrischen Satz von Wulff für die Gleichgewichtsform von Kristallen, *Z. Kristallogr., Mineral. Petrogr.* 105 (1944) 304–314, Abt. A.
- [18] X. Fan, An imbedding theorem for Musielak–Sobolev spaces, *Nonlinear Anal.* 75 (4) (2012) 1959–1971.
- [19] F. Faraci, C. Farkas, A quasilinear elliptic problem involving critical Sobolev exponents, *Collect. Math.* 66 (2) (2015) 243–259.
- [20] C. Farkas, Critical elliptic equations on non-compact Finsler manifolds, arXiv:2010.07686, 2020.
- [21] C. Farkas, J. Fodor, A. Kristály, Anisotropic elliptic problems involving sublinear terms, in: 2015 IEEE 10th Jubilee International Symposium on Applied Computational Intelligence and Informatics, Timisoara, 2015, pp. 141–146.
- [22] C. Farkas, A. Kristály, C. Varga, Singular Poisson equations on Finsler–Hadamard manifolds, *Calc. Var. Partial Differ. Equ.* 54 (2) (2015) 1219–1241.
- [23] V. Ferone, B. Kawohl, Remarks on a Finsler–Laplacian, *Proc. Am. Math. Soc.* 137 (1) (2009) 247–253.
- [24] L. Gasiński, N.S. Papageorgiou, Constant sign and nodal solutions for superlinear double phase problems, *Adv. Calc. Var.*, <https://doi.org/10.1515/acv-2019-0040>.
- [25] L. Gasiński, P. Winkert, Constant sign solutions for double phase problems with superlinear nonlinearity, *Nonlinear Anal.* 195 (2020) 111739.
- [26] L. Gasiński, P. Winkert, Existence and uniqueness results for double phase problems with convection term, *J. Differ. Equ.* 268 (8) (2020) 4183–4193.

- [27] L. Gasiński, P. Winkert, Sign changing solution for a double phase problem with nonlinear boundary condition via the Nehari manifold, *J. Differ. Equ.* 274 (2021) 1037–1066.
- [28] A.V. Lair, A.W. Shaker, Classical and weak solutions of a singular semilinear elliptic problem, *J. Math. Anal. Appl.* 211 (2) (1997) 193–222.
- [29] P. Lindqvist, On the equation $\operatorname{div}(|\nabla u|^{p-2}\nabla u) + \lambda|u|^{p-2}u = 0$, *Proc. Am. Math. Soc.* 109 (1) (1990) 157–164.
- [30] W. Liu, G. Dai, Existence and multiplicity results for double phase problem, *J. Differ. Equ.* 265 (9) (2018) 4311–4334.
- [31] P. Marcellini, Regularity and existence of solutions of elliptic equations with p, q -growth conditions, *J. Differ. Equ.* 90 (1) (1991) 1–30.
- [32] P. Marcellini, The stored-energy for some discontinuous deformations in nonlinear elasticity, in: *Partial Differential Equations and the Calculus of Variations*, Vol. II, vol. 2, Birkhäuser Boston, Boston, 1989, pp. 767–786.
- [33] G. Marino, P. Winkert, Existence and uniqueness of elliptic systems with double phase operators and convection terms, *J. Math. Anal. Appl.* 492 (2020) 124423.
- [34] S.-I. Ohta, Uniform convexity and smoothness, and their applications in Finsler geometry, *Math. Ann.* 343 (3) (2009) 669–699.
- [35] J. Ok, Partial regularity for general systems of double phase type with continuous coefficients, *Nonlinear Anal.* 177 (2018) 673–698.
- [36] J. Ok, Regularity for double phase problems under additional integrability assumptions, *Nonlinear Anal.* 194 (2020) 111408.
- [37] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, Ground state and nodal solutions for a class of double phase problems, *Z. Angew. Math. Phys.* 71 (1) (2020), 15 pp.
- [38] N.S. Papageorgiou, P. Winkert, *Applied Nonlinear Functional Analysis. An Introduction*, De Gruyter, Berlin, 2018.
- [39] N.S. Papageorgiou, P. Winkert, Singular Dirichlet (p, q) -equations, *Mediterr. J. Math.* (2021), in press.
- [40] N.S. Papageorgiou, P. Winkert, Singular p -Laplacian equations with superlinear perturbation, *J. Differ. Equ.* 266 (2–3) (2019) 1462–1487.
- [41] K. Perera, M. Squassina, Existence results for double-phase problems via Morse theory, *Commun. Contemp. Math.* 20 (2) (2018) 1750023.
- [42] M.A. Ragusa, A. Tachikawa, Regularity for minimizers for functionals of double phase with variable exponents, *Adv. Nonlinear Anal.* 9 (1) (2020) 710–728.
- [43] G. Randers, On an asymmetrical metric in the four-space of general relativity, *Phys. Rev.* 59 (2) (1941) 195–199.
- [44] Y. Sun, S. Wu, Y. Long, Combined effects of singular and superlinear nonlinearities in some singular boundary value problems, *J. Differ. Equ.* 176 (2) (2001) 511–531.
- [45] J.E. Taylor, Crystalline variational methods, *Proc. Natl. Acad. Sci. USA* 99 (24) (2002) 15277–15280.
- [46] G. Wang, C. Xia, A characterization of the Wulff shape by an overdetermined anisotropic PDE, *Arch. Ration. Mech. Anal.* 199 (1) (2011) 99–115.
- [47] S.D. Zeng, Y.R. Bai, L. Gasiński, P. Winkert, Existence results for double phase implicit obstacle problems involving multivalued operators, *Calc. Var. Partial Differ. Equ.* 59 (5) (2020) 176.
- [48] S.D. Zeng, L. Gasiński, P. Winkert, Y.R. Bai, Existence of solutions for double phase obstacle problems with multivalued convection term, *J. Math. Anal. Appl.* (2021), <https://doi.org/10.1016/j.jmaa.2020.123997>, in press.
- [49] V.V. Zhikov, Averaging of functionals of the calculus of variations and elasticity theory, *Izv. Akad. Nauk SSSR, Ser. Mat.* 50 (4) (1986) 675–710.
- [50] V.V. Zhikov, S.M. Kozlov, O.A. Oleĭnik, *Homogenization of Differential Operators and Integral Functionals*, Springer-Verlag, Berlin, 1994.