



(p, q) -Equations with Negative Concave Terms

Ángel Crespo-Blanco¹ · Nikolaos S. Papageorgiou² · Patrick Winkert¹

Received: 6 May 2022 / Accepted: 17 August 2022
© The Author(s) 2022

Abstract

In this paper, we study a nonlinear Dirichlet problem driven by the (p, q) -Laplacian and with a reaction that has the combined effects of a negative concave term and of an asymmetric perturbation which is superlinear on the positive semiaxis and resonant in the negative one. We prove a multiplicity theorem for such problems obtaining three nontrivial solutions, all with sign information. Furthermore, under a local symmetry condition, we prove the existence of a whole sequence of sign-changing solutions converging to zero in $C_0^1(\bar{\Omega})$.

Keywords Concave and convex nonlinearities · Constant sign and nodal solutions · Critical groups · (p, q) -Laplacian · Regularity theory · Resonance

Mathematics Subject Classification 35J20 · 35J60 · 58E05

1 Introduction

Let $\Omega \subseteq \mathbb{R}^N$ be a bounded domain with a C^2 -boundary $\partial\Omega$. In this paper, we study the following nonlinear Dirichlet problem

$$\begin{aligned} -\Delta_p u - \Delta_q u &= \vartheta(x)|u|^{\tau-2}u + f(x, u) \quad \text{in } \Omega, \\ u|_{\partial\Omega} &= 0, \quad 1 < \tau < q < p, \end{aligned} \tag{1.1}$$

✉ Patrick Winkert
winkert@math.tu-berlin.de

Ángel Crespo-Blanco
crespo@math.tu-berlin.de

Nikolaos S. Papageorgiou
npapg@math.ntua.gr

¹ Institut für Mathematik, Technische Universität Berlin, Straße des 17. Juni 136, 10623 Berlin, Germany

² Department of Mathematics, National Technical University, Zografou Campus, 15780 Athens, Greece

where Δ_r denotes the r -Laplacian for $r \in (1, \infty)$ given by

$$\Delta_r u = \operatorname{div} (|\nabla u|^{r-2} \nabla u) \quad \text{for } u \in W_0^{1,r}(\Omega).$$

Problem (1.1) is driven by the sum of two such operators with different exponents called the (p, q) -Laplacian which is a nonhomogeneous operator. For such problems, we refer to the survey paper of Marano and Mosconi [13] and the references therein. In the right-hand side of (1.1), we have the combined effects of two distinct nonlinear terms. One term is the power function $s \rightarrow \vartheta(x)|s|^{\tau-2}s$ with $1 < \tau < q$ and $0 > -c_0 \geq \vartheta(\cdot) \in L^\infty(\Omega)$ which is a concave contribution (so $(q-1)$ -sublinear) to the reaction. The perturbation $f: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function, that is, $x \rightarrow f(x, s)$ is measurable for all $s \in \mathbb{R}$ and $s \rightarrow f(x, s)$ is continuous for a.a. $x \in \Omega$, which exhibits asymmetric growth as $s \rightarrow \pm\infty$. To be more precise, $f(x, \cdot)$ is $(p-1)$ -linear in the negative semiaxis (as $s \rightarrow -\infty$) and can be resonant with respect to the principal eigenvalue of $(-\Delta_p, W_0^{1,p}(\Omega))$. In the positive semiaxis (as $s \rightarrow +\infty$), $f(x, \cdot)$ is $(p-1)$ -superlinear but without satisfying the Ambrosetti–Rabinowitz condition (AR-condition for short). Hence, problem (1.1) is partly resonant and partly a concave–convex problem. In addition to this lack of symmetric behavior, another feature which distinguishes our work here from earlier ones on nonlinear elliptic equations with concave terms, is the fact that the coefficient $\vartheta: \Omega \rightarrow \mathbb{R}$ of the concave term is x -dependent and negative. In the past, problems with a negative concave term were studied by Perera [22], de Paiva and Massa [3], Papageorgiou et al. [20] for semilinear equations and by Papageorgiou and Winkert [15] for nonlinear equations driven by the $(p, 2)$ -Laplacian. From these works only the paper of Papageorgiou et al. [20] considers perturbations with asymmetric behavior as $s \rightarrow \pm\infty$. In the literature, papers dealing with equations with concave terms assume that the coefficient is a positive constant. This is the case in the classical concave–convex problems, see Ambrosetti et al. [2] for equations driven by the Laplacian and by García Azorero et al. [5] for equations driven by the p -Laplacian. The difficulty that we encounter when we deal with equations that have negative concave terms is that the nonlinear strong maximum principle is not applicable, see Pucci and Serrin [23].

2 Preliminaries

In this section, we will recall the basic facts about the function spaces, the properties of the operator and some results of Morse theory.

To this end, let $\Omega \subseteq \mathbb{R}^N$ be a bounded domain with a C^2 -boundary $\partial\Omega$. For any $r \in [1, \infty]$, we denote by $L^r(\Omega) = L^r(\Omega; \mathbb{R})$ and $L^r(\Omega; \mathbb{R}^N)$ the usual Lebesgue spaces with the norm $\|\cdot\|_r$. Moreover, the Sobolev space $W_0^{1,r}(\Omega)$ is equipped with the equivalent norm $\|\cdot\| = \|\nabla \cdot\|_r$ for $1 < r < \infty$.

The Banach space

$$C_0^1(\overline{\Omega}) = \left\{ u \in C^1(\overline{\Omega}) : u|_{\partial\Omega} = 0 \right\}$$

is an ordered Banach space with positive cone

$$C_0^1(\bar{\Omega})_+ = \left\{ u \in C_0^1(\bar{\Omega}) : u(x) \geq 0 \text{ for all } x \in \bar{\Omega} \right\}.$$

This cone has a nonempty interior given by

$$\text{int} \left(C_0^1(\bar{\Omega})_+ \right) = \left\{ u \in C_0^1(\bar{\Omega})_+ : u(x) > 0 \text{ for all } x \in \Omega, \frac{\partial u}{\partial n}(x) < 0 \text{ for all } x \in \partial\Omega \right\},$$

where $n(\cdot)$ stands for the outward unit normal on $\partial\Omega$.

For $r \in (1, \infty)$, we denote by $\hat{\lambda}_1(r)$, the first eigenvalue of $(-\Delta_r, W_0^{1,r}(\Omega))$. We know that $\hat{\lambda}_1(r) > 0$ and

$$\hat{\lambda}_1(r) = \inf_{u \in W_0^{1,r}(\Omega) \setminus \{0\}} \frac{\|\nabla u\|_r^r}{\|u\|_r^r}. \tag{2.1}$$

Furthermore, $\hat{\lambda}_1(r)$ is isolated, simple, and the infimum in (2.1) is achieved on the corresponding one-dimensional eigenspace, see L \hat{e} [10]. The elements of this eigenspace have fixed sign. By $\hat{u}_1(r)$, we denote the positive, L^r -normalized (that is, $\|\hat{u}_1(r)\|_r = 1$) eigenfunction related to $\hat{\lambda}_1(r)$. The nonlinear regularity theory and the nonlinear Hopf maximum principle imply that $\hat{u}_1(r) \in \text{int} \left(C_0^1(\bar{\Omega})_+ \right)$.

We also use the weighted eigenvalue problem

$$\begin{aligned} -\Delta_p u &= \tilde{\lambda} \xi(x) |u|^{p-2} u && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega, \end{aligned} \tag{2.2}$$

with eigenvalue $\tilde{\lambda} > 0$ and $\xi \in L^\infty(\Omega)_+ \setminus \{0\}$. We know that if $\xi_1(x) \leq \xi_2(x)$ a. e. in Ω and $\xi_1 \neq \xi_2$, then $\tilde{\lambda}_1(p, \xi_2) < \tilde{\lambda}_1(p, \xi_1)$, see Motreanu et al. [14, Proposition 9.47(d)].

Let $A_r : W_0^{1,r}(\Omega) \rightarrow W^{-1,r'}(\Omega) = W_0^{1,r}(\Omega)^*$ with $\frac{1}{r} + \frac{1}{r'} = 1$ be the nonlinear operator defined by

$$\langle A_r(u), h \rangle = \int_{\Omega} |\nabla u|^{r-2} \nabla u \cdot \nabla h \, dx \quad \text{for all } u, h \in W_0^{1,r}(\Omega),$$

where $\langle \cdot, \cdot \rangle$ is the duality pairing between $W_0^{1,r}(\Omega)$ and its dual space $W_0^{1,r}(\Omega)^*$. This operator is bounded, continuous, strictly monotone, and of type (S_+) , that is,

$$u_n \rightharpoonup u \text{ in } W_0^{1,r}(\Omega) \quad \text{and} \quad \limsup_{n \rightarrow \infty} \langle A_r(u_n), u_n - u \rangle \leq 0$$

imply $u_n \rightarrow u$ in $W_0^{1,r}(\Omega)$, see Motreanu et al. [14, p.40].

Let X be a Banach space, $\varphi \in C^1(X)$ and $c \in \mathbb{R}$. We introduce the following two sets

$$K_\varphi = \{u \in X : \varphi'(u) = 0\} \quad \text{and} \quad \varphi^c = \{u \in X : \varphi(u) \leq c\}.$$

If (Y_1, Y_2) is a topological pair such that $Y_2 \subseteq Y_1 \subset X$ and $k \in \mathbb{N}_0$, then we denote by $H_k(Y_1, Y_2)$ the k -th singular homology group for the pair (Y_1, Y_2) with integer coefficients. If $u \in K_\varphi$ is isolated, the k -th critical group of φ at u is defined by

$$C_k(\varphi, u) = H_k(\varphi^c \cap U, \varphi^c \cap U \setminus \{u\}), \quad k \in \mathbb{N}_0,$$

with $c = \varphi(u)$ and U being an open neighborhood of u such that $\varphi^c \cap K_\varphi \cap U = \{u\}$. The excision property of singular homology implies that the definition of $C_k(\varphi, u)$ is independent of the choice of the isolating neighborhood U , see Motreanu et al. [14]. The usage of critical groups allows us to distinguish between critical points of the energy functional.

We say that $\varphi \in C^1(X)$ satisfies the Cerami condition (C-condition for short) if every sequence $\{u_n\}_{n \in \mathbb{N}} \subseteq X$ such that $\{\varphi(u_n)\}_{n \in \mathbb{N}} \subseteq \mathbb{R}$ is bounded and $(1 + \|u_n\|_X)\varphi'(u_n) \rightarrow 0$ in X^* has a strongly convergent subsequence. This is a compactness-type condition on the functional φ which compensates the fact that the ambient space X need not be locally compact.

For $s \in \mathbb{R}$, we set $s^\pm = \max\{\pm s, 0\}$. If $u: \Omega \rightarrow \mathbb{R}$ is a measurable function, we define $u^\pm(x) = u(x)^\pm$ for all $x \in \Omega$. If $u \in W_0^{1,p}(\Omega)$, then $u^\pm \in W_0^{1,p}(\Omega)$ and $u = u^+ - u^-$ as well as $|u| = u^+ + u^-$. If $u, v: \Omega \rightarrow \mathbb{R}$ are two measurable functions such that $u(x) \leq v(x)$ for all $x \in \Omega$, then we define

$$[u, v] = \left\{ h \in W_0^{1,p}(\Omega) : u(x) \leq h(x) \leq v(x) \text{ for a. a. } x \in \Omega \right\}.$$

Moreover, we denote by $\text{int}_{C_0^1(\overline{\Omega})}[u, v]$ the interior of $[u, v] \cap C_0^1(\overline{\Omega})$ in $C_0^1(\overline{\Omega})$. Finally, the critical Sobolev exponent of $p \in (1, \infty)$, denoted by p^* , is given by

$$p^* = \begin{cases} \frac{Np}{N-p} & \text{if } p < N, \\ +\infty & \text{if } N \leq p. \end{cases}$$

3 Multiple Solutions

In this section, we produce three nontrivial solutions of problem (1.1) where two of them have constant sign and one has changing sign.

Now we introduce the hypotheses on the data of problem (1.1).

H_0 : $\vartheta \in L^\infty(\Omega)$ and $\vartheta(x) \leq -c_0 < 0$ for a. a. $x \in \Omega$.

Remark 3.1 It is an interesting open question if the results in this paper remain valid under the weaker condition $\vartheta(x) < 0$ for a. a. $x \in \Omega$.

H_1 : $f: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function such that $f(x, 0) = 0$ for a. a. $x \in \Omega$ and it satisfies the following assumptions:

(i) there exist $r \in (p, p^*)$ and $0 \leq a(\cdot) \in L^\infty(\Omega)$ such that

$$|f(x, s)| \leq a(x) \left(1 + |s|^{r-1}\right)$$

for a. a. $x \in \Omega$ and for all $s \in \mathbb{R}$;

(ii) if $F(x, s) = \int_0^s f(x, t) dt$, then

$$\lim_{s \rightarrow +\infty} \frac{F(x, s)}{s^p} = +\infty$$

uniformly for a. a. $x \in \Omega$ and there exists

$$\mu \in \left((r - p) \max \left\{ \frac{N}{p}, 1 \right\}, p^* \right)$$

such that

$$0 < \beta_0 \leq \liminf_{s \rightarrow +\infty} \frac{f(x, s)s - pF(x, s)}{s^\mu}$$

uniformly for a. a. $x \in \Omega$;

(iii) there exist $\beta_1 \in L^\infty(\Omega)$ and $\beta_2 > 0$ such that

$$\hat{\lambda}_1(p) \leq \beta_1(x) \quad \text{for a. a. } x \in \Omega$$

with $\beta_1 \not\equiv \hat{\lambda}_1(p)$ and

$$\beta_1(x) \leq \liminf_{s \rightarrow -\infty} \frac{f(x, s)}{|s|^{p-2}s} \leq \limsup_{s \rightarrow -\infty} \frac{f(x, s)}{|s|^{p-2}s} \leq \beta_2$$

uniformly for a. a. $x \in \Omega$.

(iv) there exists $\beta \in (1, \tau)$ such that

$$\lim_{s \rightarrow 0} \frac{f(x, s)}{|s|^{\beta-2}s} = 0$$

uniformly for a. a. $x \in \Omega$,

$$\liminf_{s \rightarrow 0} \frac{f(x, s)}{|s|^{\tau-2}s} \geq \eta > \|\vartheta\|_\infty$$

uniformly for a. a. $x \in \Omega$ and for every $\lambda > 0$ there exists $\hat{\mu}(\lambda) \in (1, \beta)$ such that $\hat{\mu}(\lambda) \rightarrow \hat{\mu} \in (1, \beta)$ as $\lambda \rightarrow 0^+$ and

$$f(x, s)s \leq \hat{c} \left(\lambda |s|^{\hat{\mu}(\lambda)} + |s|^r \right) - \tilde{c} |s|^\beta$$

for a. a. $x \in \Omega$, for all $s \in \mathbb{R}$ with $\hat{c}, \tilde{c} > 0$.

Remark 3.2 Hypotheses H_1 (ii) and H_1 (iii) imply the asymmetric behavior of the perturbation $f(x, \cdot)$. Indeed, hypothesis H_1 (ii) says that $f(x, \cdot)$ is $(p - 1)$ -superlinear as $s \rightarrow +\infty$ but need not satisfy the AR-condition, see, for example, Ghoussoub [6, p. 59]. Our condition is less restrictive and allows also nonlinearities with “slower” growth as $s \rightarrow +\infty$ which fail to satisfy the AR-condition. Here, we refer to a unilateral version of the condition since it concerns only the positive semiaxis $[0, \infty)$. Hypothesis H_1 (iii) says that $f(x, \cdot)$ is $(p - 1)$ -linear as $s \rightarrow -\infty$ and can be resonant with respect to the principal eigenvalue of $(-\Delta_p, W_0^{1,p}(\Omega))$. Note that in hypothesis H_1 (i), we want $a \in L^\infty(\Omega)$ in order to be able to apply the regularity theory of Lieberman [12].

Example 3.3 The following function satisfies hypotheses H_1 but fails to satisfy the AR-condition:

$$f(x, s) = \begin{cases} \gamma(x) (|s|^{p-2}s - |s|^{q-2}s) & \text{if } s < -1, \\ \eta(x) (|s|^{\tau-2}s - |s|^{\mu-2}s) & \text{if } -1 \leq s \leq 1, \\ cs^{p-1} \ln(s) & \text{if } 1 < s, \end{cases}$$

with $\gamma \in L^\infty(\Omega)$, $\gamma(x) \geq \hat{\lambda}_1(q)$, $\gamma \not\equiv \hat{\lambda}_1(q)$ and $\eta \in L^\infty(\Omega)$, $\text{ess inf}_\Omega \eta > \|\vartheta\|_\infty$, $c > 0$ and $p > \mu > \tau$.

Let $\varphi: W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ be the energy functional corresponding to problem (1.1) defined by

$$\varphi(u) = \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \frac{1}{\tau} \int_\Omega \vartheta(x) |u|^\tau \, dx - \int_\Omega F(x, u) \, dx$$

for all $u \in W_0^{1,p}(\Omega)$. It is clear that $\varphi \in C^1(W_0^{1,p}(\Omega))$. Moreover, we introduce the positive and negative truncations of φ , namely, the C^1 -functionals $\varphi_\pm: W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ given by

$$\varphi_\pm(u) = \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \frac{1}{\tau} \int_\Omega \vartheta(x) (u^\pm)^\tau \, dx - \int_\Omega F(x, \pm u^\pm) \, dx$$

for all $u \in W_0^{1,p}(\Omega)$.

Our idea is to work with the truncated functionals $\varphi_\pm: W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$.

Proposition 3.4 *Let hypotheses H_0 and H_1 be satisfied. Then there exists $\hat{q} > 0$ such that*

$$\varphi_\pm(u) \geq m > 0 \text{ for all } u \in W_0^{1,p}(\Omega) \text{ with } \|u\| = \hat{q}.$$

Proof From hypotheses H_1 (iv), we see that for given $\varepsilon > 0$, we can find $c_1 = c_1(\varepsilon) > 0$ such that

$$F(x, s) \leq \frac{\varepsilon - \tilde{c}}{\beta} |s|^\beta + c_1 \left(\lambda |s|^{\hat{\mu}(\lambda)} + |s|^r \right) \quad \text{for a. a. } x \in \Omega \text{ and for all } s \in \mathbb{R}. \tag{3.1}$$

Using (3.1) and hypotheses H_0 , we get for $u \in W_0^{1,p}(\Omega)$

$$\varphi_\pm(u) \geq \left(\frac{1}{p} - \lambda c_2 \|u\|^{\hat{\mu}(\lambda)-p} - c_3 \|u\|^{r-p} \right) \|u\|^p$$

for some $c_2, c_3 > 0$.

Let

$$\xi_\lambda(t) = \lambda c_2 t^{\hat{\mu}(\lambda)-p} + c_3 t^{r-p} \quad \text{for } t > 0.$$

Since $\hat{\mu}(\lambda) < \beta < p < r$, we see that

$$\xi_\lambda(t) \rightarrow +\infty \quad \text{as } t \rightarrow 0^+ \text{ and as } t \rightarrow +\infty.$$

Therefore, we find a number $t_0 \in (0, \infty)$ such that

$$\xi_\lambda(t_0) = \inf_{t>0} \xi_\lambda(t).$$

Thus, $\xi'_\lambda(t_0) = 0$, and this implies

$$t_0 = \left[\frac{\lambda c_2 (p - \hat{\mu}(\lambda))}{c_3 (r - p)} \right]^{\frac{1}{r - \hat{\mu}(\lambda)}}.$$

Since $\xi_\lambda(t_0) \rightarrow 0$ as $\lambda \rightarrow 0^+$, there exists $\lambda_0 > 0$ such that

$$\xi_\lambda(t_0) < \frac{1}{p} \quad \text{for all } \lambda \in (0, \lambda_0).$$

Fix $\lambda \in (0, \lambda_0)$, then, for $\|u\| = t_0$, we have

$$\varphi_\pm(u) > 0.$$

Next, we show that $\varphi_+ : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ satisfies the C-condition.

Proposition 3.5 *Let hypotheses H_0 and H_1 be satisfied. Then the functional $\varphi_+ : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ satisfies the C-condition.*

Proof Let $\{u_n\}_{n \in \mathbb{N}} \subseteq W_0^{1,p}(\Omega)$ be a sequence such that

$$|\varphi_+(u_n)| \leq c_3 \quad \text{for some } c_3 > 0 \text{ and for all } n \in \mathbb{N}, \tag{3.2}$$

$$(1 + \|u_n\|) \varphi'_+(u_n) \rightarrow 0 \quad \text{in } W^{-1,p'}(\Omega). \tag{3.3}$$

From (3.3), we get

$$\begin{aligned} & \left| \langle A_p(u_n), h \rangle + \langle A_q(u_n), h \rangle - \int_{\Omega} \vartheta(x) (u_n^+)^{\tau-1} h \, dx - \int_{\Omega} f(x, u_n^+) h \, dx \right| \\ & \leq \frac{\varepsilon_n \|h\|}{1 + \|u_n\|} \quad \text{for all } h \in W_0^{1,p}(\Omega) \text{ with } \varepsilon_n \rightarrow 0^+. \end{aligned} \tag{3.4}$$

Choosing $h = -u_n^- \in W_0^{1,p}(\Omega)$ in (3.4) gives $\|u_n^-\|^p \leq \varepsilon_n$ for all $n \in \mathbb{N}$ and so

$$u_n^- \rightarrow 0 \quad \text{in } W_0^{1,p}(\Omega). \tag{3.5}$$

Combining (3.2) and (3.5) yields

$$\|\nabla u_n^+\|_p^p + \frac{p}{q} \|\nabla u_n^+\|_q^q - \frac{p}{\tau} \int_{\Omega} \vartheta(x) (u_n^+)^{\tau} \, dx - \int_{\Omega} pF(x, u_n^+) \, dx \leq c_4 \tag{3.6}$$

for some $c_4 > 0$ and for all $n \in \mathbb{N}$. Next, we take $h = u_n^+ \in W_0^{1,p}(\Omega)$ in (3.4). We obtain

$$-\|\nabla u_n^+\|_p^p - \|\nabla u_n^+\|_q^q + \int_{\Omega} \vartheta(x) (u_n^+)^{\tau} \, dx + \int_{\Omega} f(x, u_n^+) u_n^+ \, dx \leq \varepsilon_n \tag{3.7}$$

for all $n \in \mathbb{N}$. Adding (3.6) and (3.7) and using hypotheses H_0 as well as $\tau < q < p$, we get

$$\int_{\Omega} (f(x, u_n^+) u_n^+ - pF(x, u_n^+)) \, dx \leq c_5 \tag{3.8}$$

for some $c_5 > 0$ and for all $n \in \mathbb{N}$.

Hypotheses H_1 (i) and H_1 (ii) imply that we can find $\hat{\beta}_0 \in (0, \beta_0)$ and $c_6 > 0$ such that

$$\hat{\beta}_0 s^{\mu} - c_6 \leq f(x, s) s - pF(x, s) \tag{3.9}$$

for a.a. $x \in \Omega$ and for all $s \geq 0$. Using (3.9) in (3.8) leads to

$$\|u_n^+\|_{\mu}^{\mu} \leq c_7 \quad \text{for some } c_7 > 0 \text{ and for all } n \in \mathbb{N}.$$

Hence

$$\{u_n^+\}_{n \in \mathbb{N}} \subseteq L^{\mu}(\Omega) \text{ is bounded.} \tag{3.10}$$

First, assume that $p \neq N$. From hypothesis H_1 (ii) it is clear that we may assume that $\mu < r < p^*$. Then we can find $t \in (0, 1)$ such that

$$\frac{1}{r} = \frac{1-t}{\mu} + \frac{t}{p^*}. \tag{3.11}$$

Using the interpolation inequality (see Papageorgiou and Winkert [18, p. 116]), we have

$$\|u_n^+\|_r \leq \|u_n^+\|_\mu^{1-t} \|u_n^+\|_{p^*}^t \quad \text{for all } n \in \mathbb{N}.$$

This combined with (3.10) results in

$$\|u_n^+\|_r^r \leq c_8 \|u_n^+\|^{tr} \quad \text{for all } n \in \mathbb{N} \tag{3.12}$$

with some $c_8 > 0$. Testing (3.4) with $h = u_n^+ \in W_0^{1,p}(\Omega)$ we obtain

$$\|\nabla u_n^+\|_p^p \leq \varepsilon_n + \int_\Omega f(x, u_n^+) u_n^+ dx \quad \text{for all } n \in \mathbb{N}$$

due to hypotheses H_0 . Using H_1 (i), this implies

$$\|u_n^+\|^p \leq c_9 (1 + \|u_n^+\|_r^r) \quad \text{for all } n \in \mathbb{N}$$

with some $c_9 > 0$. Combining this with (3.12) yields

$$\|u_n^+\|^p \leq c_{10} (1 + \|u_n^+\|^{tr}) \quad \text{for all } n \in \mathbb{N} \tag{3.13}$$

for some $c_{10} > 0$.

Recall that $p \neq N$. If $p > N$, then by definition we have $p^* = \infty$ and so

$$\frac{1}{r} = \frac{1-t}{\mu},$$

see (3.11), which implies, because of H_1 (ii), that $tr = r - \mu < p$. Then we conclude from (3.13) that

$$\{u_n^+\}_{n \in \mathbb{N}} \subseteq W_0^{1,p}(\Omega) \text{ is bounded.} \tag{3.14}$$

If $p < N$, then we have by definition $p^* = \frac{Np}{N-p}$. So from (3.11) and H_1 (ii), it follows

$$tr = \frac{p^*(r - \mu)}{p^* - \mu} = \frac{Np(r - \mu)}{Np - N\mu + \mu p} < \frac{Np(r - \mu)}{Np - N\mu + (r - p)\frac{N}{p}p} = p.$$

Hence, (3.14) holds again in this case.

Finally, let $p = N$. Then by the Sobolev embedding theorem, we know that $W_0^{1,p}(\Omega) \hookrightarrow L^s(\Omega)$ is continuous for all $1 \leq s < \infty$. Then, in the argument above, we need to replace p^* by $s > r > \mu$. We choose $t \in (0, 1)$ such that

$$\begin{aligned} \text{which gives} \quad \frac{1}{r} &= \frac{1-t}{\mu} + \frac{t}{s}, \\ tr &= \frac{s(r-\mu)}{s-\mu}. \end{aligned} \quad (3.15)$$

Note that $\frac{s(r-\mu)}{s-\mu} \rightarrow r - \mu$ as $s \rightarrow +\infty$ and $r - \mu < p$, see H₁ (ii). We choose $s > r$ large enough such that

$$\frac{s(r-\mu)}{s-\mu} < p.$$

Then, using (3.15), we have $tr < p$ and so $\{u_n^+\}_{n \in \mathbb{N}} \subseteq W_0^{1,p}(\Omega)$ is bounded. Combining this with (3.5), we obtain that $\{u_n\}_{n \in \mathbb{N}} \subseteq W_0^{1,p}(\Omega)$ is bounded.

Then there exists a subsequence, not relabeled, such that

$$u_n \rightarrow u \quad \text{in } W_0^{1,p}(\Omega) \quad \text{and} \quad u_n \rightarrow u \quad \text{in } L^r(\Omega). \quad (3.16)$$

If we use $h = u_n - u \in W_0^{1,p}(\Omega)$ in (3.4), pass to the limit as $n \rightarrow \infty$ and use (3.16), we obtain

$$\lim_{n \rightarrow \infty} (\langle A_p(u_n), u_n - u \rangle + \langle A_q(u_n), u_n - u \rangle) = 0.$$

By the monotonicity of A_q , we have

$$\langle A_q(u), u_n - u \rangle \leq \langle A_q(u_n), u_n - u \rangle.$$

Using this in the limit above, we obtain

$$\limsup_{n \rightarrow \infty} (\langle A_p(u_n), u_n - u \rangle + \langle A_q(u), u_n - u \rangle) \leq 0.$$

Hence, from the convergence properties in (3.16), we conclude that

$$\limsup_{n \rightarrow \infty} \langle A_p(u_n), u_n - u \rangle \leq 0.$$

The (S_+) -property of A_p implies that $u_n \rightarrow u$ in $W_0^{1,p}(\Omega)$. This shows that φ_+ satisfies the C-condition.

Proposition 3.5 leads to the following existence result for problem (1.1).

Proposition 3.6 *Let hypotheses H_0 and H_1 . Then problem (1.1) has at least one positive solution $u_0 \in C_0^1(\overline{\Omega})_+ \setminus \{0\}$.*

Proof From Proposition 3.4, we know that

$$\varphi_+(0) = 0 < m \leq \varphi_+(u) \quad \text{for all } u \in W_0^{1,p}(\Omega) \text{ with } \|u\| = \hat{\varrho}. \tag{3.17}$$

Also, from Proposition 3.5, we know that

$$\varphi_+ : W_0^{1,p}(\Omega) \rightarrow \mathbb{R} \text{ satisfies the C-condition.} \tag{3.18}$$

Moreover, hypothesis H_1 (ii) implies that if $u \in \text{int}(C_0^1(\overline{\Omega})_+)$, then

$$\varphi_+(tu) \rightarrow -\infty \quad \text{as } t \rightarrow +\infty. \tag{3.19}$$

Then, (3.17), (3.18), and (3.19) permit the usage of the mountain pass theorem. Therefore, we can find $u_0 \in W_0^{1,p}(\Omega)$ such that

$$u_0 \in K_{\varphi_+} \quad \text{and} \quad \varphi_+(0) = 0 < m \leq \varphi_+(u_0).$$

Hence, $u_0 \neq 0$. From Ho et al. [7, Theorem 3.1], we know that $u_0 \in L^\infty(\Omega)$. Then the nonlinear regularity theory of Lieberman [12] implies that $u_0 \in C_0^1(\overline{\Omega})_+ \setminus \{0\}$.

Remark 3.7 Eventually, we will show that $u_0 \in \text{int}(C_0^1(\overline{\Omega})_+)$, see Corollary 3.12. However, at this point, due to the negative concave term, we cannot use the nonlinear Hopf maximum principle, see Pucci and Serrin [23, p.120], and infer that $u_0 \in \text{int}(C_0^1(\overline{\Omega})_+)$.

Next, we are looking for a negative solution of problem (1.1). So, we work with the functional $\varphi_- : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$. For the functional $\varphi_- : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$, we have the following proposition.

Proposition 3.8 *Let hypotheses H_0 and H_1 be satisfied. Then the functional $\varphi_- : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ satisfies the C-condition.*

Proof Let $\{u_n\}_{n \in \mathbb{N}} \subseteq W_0^{1,p}(\Omega)$ be a sequence such that $\{\varphi_-(u_n)\}_{n \in \mathbb{N}} \subseteq \mathbb{R}$ is bounded and

$$(1 + \|u_n\|) \varphi'_-(u_n) \rightarrow 0 \quad \text{in } W^{-1,p'}(\Omega). \tag{3.20}$$

From (3.20), we have

$$\begin{aligned} & \left| \langle A_p(u_n), h \rangle + \langle A_q(u_n), h \rangle + \int_{\Omega} \vartheta(x) (u_n^-)^{\tau-1} h \, dx - \int_{\Omega} f(x, -u_n^-) h \, dx \right| \\ & \leq \frac{\varepsilon_n \|h\|}{1 + \|u_n\|} \quad \text{for all } h \in W_0^{1,p}(\Omega) \text{ with } \varepsilon_n \rightarrow 0^+. \end{aligned} \tag{3.21}$$

If we choose $h = u_n^+ \in W_0^{1,p}(\Omega)$ in (3.21), we obtain $\|u_n^+\|^p \leq \varepsilon_n$ for all $n \in \mathbb{N}$ which implies

$$u_n^+ \rightarrow 0 \text{ in } W_0^{1,p}(\Omega). \tag{3.22}$$

Suppose that $\|u_n^-\| \rightarrow \infty$ and let $y_n = \frac{u_n^-}{\|u_n^-\|}$. Then $\|y_n\| = 1$ for all $n \in \mathbb{N}$. Therefore, we may suppose, for a subsequence if necessary, that

$$y_n \rightharpoonup y \text{ in } W_0^{1,p}(\Omega) \text{ and } y_n \rightarrow y \text{ in } L^p(\Omega) \tag{3.23}$$

for some $y \in W_0^{1,p}(\Omega)$ with $y \geq 0$. From (3.21) and (3.22), we obtain

$$\begin{aligned} & \left| \langle A_p(-y_n), h \rangle + \frac{1}{\|u_n^-\|^{p-q}} \langle A_q(-y_n), h \rangle - \int_{\Omega} \frac{\vartheta(x)}{\|u_n^-\|^{p-\tau}} y_n^{\tau-1} h \, dx \right. \\ & \left. - \int_{\Omega} \frac{f(x, -u_n^-)}{\|u_n^-\|^{p-1}} h \, dx \right| \leq \varepsilon'_n \|h\| \text{ for all } h \in W_0^{1,p}(\Omega) \text{ with } \varepsilon'_n \rightarrow 0^+. \end{aligned} \tag{3.24}$$

Choosing $h = y_n - y \in W_0^{1,p}(\Omega)$ in (3.24), passing to the limit as $n \rightarrow \infty$ and using the convergence properties in (3.23) gives

$$\lim_{n \rightarrow \infty} \langle A_p(y_n), y_n - y \rangle = 0.$$

From the (S₊)-property of $A_p: W_0^{1,p}(\Omega) \rightarrow W^{-1,p'}(\Omega) = W_0^{1,p}(\Omega)^*$, we conclude that

$$y_n \rightarrow y \text{ in } W_0^{1,p}(\Omega) \text{ with } \|y\| = 1 \text{ and } y \geq 0. \tag{3.25}$$

Note that from hypothesis H₁ (iii), we have

$$\frac{f(\cdot, -u_n(\cdot)^-)}{\|u_n^-\|^{p-1}} \rightarrow -\hat{\beta}(x)y^{p-1} \text{ in } L^{p'}(\Omega) \tag{3.26}$$

with $\hat{\beta} \in L^\infty(\Omega)$ and $\beta_1(x) \leq \hat{\beta}(x) \leq \beta_2$ for a. a. $x \in \Omega$, see Aizicovici et al. [1, proof of Proposition 16] and Motreanu et al. [14, Proof of Theorem 11.15, p.317].

So, if we pass to the limit in (3.24) as $n \rightarrow \infty$ and use (3.25) as well as (3.26), we obtain

$$\langle A_p(-y), h \rangle = - \int_{\Omega} \hat{\beta}(x)y^{p-1} h \, dx \text{ for all } h \in W_0^{1,p}(\Omega).$$

This means that

$$-\Delta_p y = \hat{\beta}(x)y^{p-1} \text{ in } \Omega, \quad y|_{\partial\Omega} = 0.$$

From (3.25), we know that $y \neq 0$ and

$$\tilde{\lambda}_1(p, \hat{\beta}) < \tilde{\lambda}_1(p, \hat{\lambda}_1(p)) = 1, \tag{3.27}$$

see (2.2). From (3.26) and (3.27), it follows that y must be sign-changing which is a contradiction to (3.25), see also Motreanu et al. [14, Proposition 9.47(b)]. Thus, $\{u_n^-\} \subseteq W_0^{1,p}(\Omega)$ is bounded; hence, $\{u_n\} \subseteq W_0^{1,p}(\Omega)$ is bounded, see (3.22). From this as in the proof of Proposition 3.5, we conclude that $\varphi_- : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ satisfies the C-condition.

On account of hypothesis H_1 (iii), we see that

$$\varphi_-(t\hat{u}_1(p)) \rightarrow -\infty \text{ as } t \rightarrow -\infty. \tag{3.28}$$

Then (3.28), Proposition 3.8, and the mountain pass theorem lead to the following result.

Proposition 3.9 *Let hypotheses H_0 and H_1 be satisfied. Then problem (1.1) has a negative solution $v_0 \in -C_0^1(\overline{\Omega}) \setminus \{0\}$.*

In what follows \mathcal{S}_+ (resp. \mathcal{S}_-) denote the set of positive (resp. negative) solutions to (1.1). From Propositions 3.6 and 3.9, we have

$$\emptyset \neq \mathcal{S}_+ \subseteq C_0^1(\overline{\Omega})_+ \setminus \{0\} \text{ and } \emptyset \neq \mathcal{S}_- \subseteq \left(-C_0^1(\overline{\Omega})_+\right) \setminus \{0\}.$$

Next, we are going to prove that \mathcal{S}_+ has a minimal element and \mathcal{S}_- a maximal one. So we have extremal constant sign solutions, that is, there is a smallest positive solution u_* and a largest negative solution v_* . These solutions will be useful in proving the existence of a sign-changing solution. Indeed, any nontrivial solution of problem (1.1) in the order interval $[v_*, u_*]$ distinct from v_* and u_* is necessarily sign-changing.

On account of hypotheses H_1 (i) and H_1 (iv), for a given $\varepsilon > 0$, we can find $\hat{c}_1 = \hat{c}_1(\varepsilon) > 0$ such that

$$f(x, s)s \geq [\eta - \varepsilon] |s|^\tau - \hat{c}_1 |s|^r$$

for a. a. $x \in \Omega$ and for all $s \in \mathbb{R}$. This implies

$$\vartheta(x) |s|^\tau + f(x, s)s \geq [\eta - \varepsilon - \|\vartheta\|_\infty] |s|^\tau - \hat{c}_1 |s|^r$$

for a. a. $x \in \Omega$ and for all $s \in \mathbb{R}$. By hypothesis H_1 (iv), we have $\eta > \|\vartheta\|_\infty$. So, choosing $\varepsilon \in (0, \eta - \|\vartheta\|_\infty)$, we have

$$\vartheta(x) |s|^\tau + f(x, s)s \geq \hat{c}_2 |s|^\tau - \hat{c}_1 |s|^r \tag{3.29}$$

for some $\hat{c}_2 > 0$, for a.a. $x \in \Omega$ and for all $s \in \mathbb{R}$. Then, (3.29) suggests that we consider the following Dirichlet (p, q) -equation

$$\begin{aligned}
 -\Delta_p u - \Delta_q u &= \hat{c}_2 |u|^{\tau-2} u - \hat{c}_1 |u|^{r-2} u && \text{in } \Omega, \\
 u|_{\partial\Omega} &= 0, \quad 1 < \tau < q < p < r < p^*, && (3.30)
 \end{aligned}$$

Similarly to Proposition 4.1 of Papageorgiou and Winkert [17], we have the following existence and uniqueness result.

Proposition 3.10 *Problem (3.30) has a unique positive solution $\bar{u} \in \text{int}(C_0^1(\bar{\Omega})_+)$ and since problem (3.30) is odd, $\bar{v} = -\bar{u} \in -\text{int}(C_0^1(\bar{\Omega})_+)$ is the unique negative solution of (3.30).*

Proof First, we show the existence of a positive solution of problem (3.30). To this end, let $\psi_+ : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ be the C^1 -functional defined by

$$\psi_+(u) = \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q + \frac{\hat{c}_1}{r} \|u^+\|_r^r - \frac{\hat{c}_2}{\tau} \|u^+\|_\tau^\tau$$

for all $u \in W_0^{1,p}(\Omega)$. Since $\tau < q < p < r$, it is clear that $\psi_+ : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ is coercive. Also, it is sequentially weakly lower semicontinuous. Therefore, there exists $\bar{u} \in W_0^{1,p}(\Omega)$ such that

$$\psi_+(\bar{u}) = \inf \left[\psi_+(u) : u \in W_0^{1,p}(\Omega) \right]. \tag{3.31}$$

Note that if $u \in \text{int}(C_0^1(\bar{\Omega})_+)$ and $t \in (0, 1)$ small enough, then $\psi_+(tu) < 0$ since $\tau < q < p < r$ and so we have $\psi_+(\bar{u}) < 0 = \psi_+(0)$. Thus, $\bar{u} \neq 0$.

From (3.31), we have $\psi'_+(\bar{u}) = 0$, that is,

$$\langle A_p(\bar{u}), h \rangle + \langle A_q(\bar{u}), h \rangle = \hat{c}_2 \int_\Omega (\bar{u}^+)^{\tau-1} h \, dx - \hat{c}_1 \int_\Omega (\bar{u}^+)^{r-1} h \, dx$$

for all $h \in W_0^{1,p}(\Omega)$. Choosing $h = -\bar{u}^- \in W_0^{1,p}(\Omega)$ in the equality above shows that $\bar{u} \geq 0$ with $\bar{u} \neq 0$. Moreover, the nonlinear regularity theory of Lieberman [12] and the nonlinear strong maximum principle, see Pucci and Serrin [23, pp. 111 and 120], imply that $\bar{u} \in \text{int}(C_0^1(\bar{\Omega})_+)$.

Next, we show the uniqueness of this positive solution. For this purpose, we introduce the functional $j : L^1(\Omega) \rightarrow \mathbb{R} \cup \{\infty\}$ defined by

$$j(u) = \begin{cases} \frac{1}{p} \|\nabla u^{\frac{1}{p}}\|_p^p + \frac{1}{q} \|\nabla u^{\frac{1}{q}}\|_q^q & \text{if } u \geq 0, \quad u^{\frac{1}{\tau}} \in W_0^{1,p}(\Omega), \\ +\infty & \text{otherwise.} \end{cases}$$

Let $\text{dom } j = \{u \in L^1(\Omega) : j(u) < \infty\}$ be the effective domain of $j : L^1(\Omega) \rightarrow \mathbb{R} \cup \{\infty\}$. Using the ideas of Díaz and Saá [4] along with the fact that the function $s \mapsto s^{\frac{\hat{q}}{\tau}}$

for $\tau < \hat{\eta}$ is increasing and convex, we know that j is convex. Let $\bar{w} \in W_0^{1,p}(\Omega)$ be another positive solution of (3.30). As done before, we get $\bar{w} \in \text{int}(C_0^1(\bar{\Omega})_+)$. From l'Hospital's rule, we have

$$\frac{\bar{u}}{\bar{w}} \in L^\infty(\Omega) \quad \text{and} \quad \frac{\bar{w}}{\bar{u}} \in L^\infty(\Omega). \tag{3.32}$$

Let $h = \bar{u}^\tau - \bar{w}^\tau \in C_0^1(\bar{\Omega})$. From (3.32), we know that $\frac{\bar{w}^\tau}{\bar{u}^\tau} \leq c$ with $c > 0$ and so $-\bar{w}^\tau \geq -c\bar{u}^\tau$. Then, for $|t|$ small enough, we have

$$\bar{u}^\tau + th = (1+t)\bar{u}^\tau - t\bar{w}^\tau \geq ((1+t) - tc)\bar{u}^\tau \geq 0.$$

Clearly, $(\bar{u}^\tau + th)^{\frac{1}{\tau}} \in W_0^{1,p}(\Omega)$. Hence, $\bar{u}^\tau + th \in \text{dom } j$. Similarly, we can show that $\bar{w}^\tau + th \in \text{dom } j$.

Then the convexity of j implies that the directional derivative of j at \bar{u}^τ and at \bar{w}^τ , respectively, in the direction h exists. Moreover, using the nonlinear Green's identity, see Papageorgiou et al. [21, p.35], we have

$$j'(\bar{u}^\tau)(h) = \frac{1}{\tau} \int_\Omega \frac{-\Delta_p \bar{u} - \Delta_q \bar{u}}{\bar{u}^{\tau-1}} h \, dx = \frac{1}{\tau} \int_\Omega [\hat{c}_2 - \hat{c}_1 \bar{u}^{r-\tau}] h \, dx,$$

$$j'(\bar{w}^\tau)(h) = \frac{1}{\tau} \int_\Omega \frac{-\Delta_p \bar{w} - \Delta_q \bar{w}}{\bar{w}^{\tau-1}} h \, dx = \frac{1}{\tau} \int_\Omega [\hat{c}_2 - \hat{c}_1 \bar{w}^{r-\tau}] h \, dx.$$

The convexity of j implies the monotonicity of j' . So, we have

$$0 \leq \frac{\hat{c}_1}{\tau} \int_\Omega [\bar{w}^{r-\tau} - \bar{u}^{r-\tau}] (\bar{u}^\tau - \bar{w}^\tau) \, dx \leq 0.$$

Thus, $\bar{u} = \bar{w}$.

Since equation (3.30) is odd, $\bar{v} = -\bar{u} \in -\text{int}(C_0^1(\bar{\Omega})_+)$ is the unique negative solution of (3.30).

Proposition 3.11 *Let hypotheses H_0 and H_1 be satisfied. Then it holds $\bar{u} \leq u$ for all $u \in \mathcal{S}_+$ and $v \leq \bar{v}$ for all $v \in \mathcal{S}_-$, where \bar{u}, \bar{v} are the unique nontrivial constant sign solutions of (3.30) given in Proposition 3.10.*

Proof Let $u \in \mathcal{S}_+$ and consider the Carathéodory function $l_+ : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$l_+(x, s) = \begin{cases} \hat{c}_2 (s^+)^{\tau-1} - \hat{c}_1 (s^+)^{r-1} & \text{if } s \leq u(x), \\ \hat{c}_2 u(x)^{\tau-1} - \hat{c}_1 u(x)^{r-1} & \text{if } u(x) < s. \end{cases} \tag{3.33}$$

We set $L_+(x, s) = \int_0^s l_+(x, t) dt$ and consider the C^1 -functional $\sigma_+ : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\sigma_+(u) = \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \int_{\Omega} L_+(x, u) dx$$

for all $u \in W_0^{1,p}(\Omega)$.

From the truncation in (3.33), it is clear that $\sigma_+ : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ is coercive. Moreover, it is also sequentially weakly lower semicontinuous. So, we can find $\tilde{u} \in W_0^{1,p}(\Omega)$ such that

$$\sigma_+(\tilde{u}) = \inf \left[\sigma_+(u) : u \in W_0^{1,p}(\Omega) \right]. \tag{3.34}$$

Since $\tau < q < p < r$, we see that $\sigma_+(\tilde{u}) < 0 = \sigma_+(0)$. Hence, $\tilde{u} \neq 0$.

From (3.34), we have $\sigma'_+(\tilde{u}) = 0$. This gives

$$\langle A_p(\tilde{u}), h \rangle + \langle A_q(\tilde{u}), h \rangle = \int_{\Omega} l_+(x, \tilde{u}) h dx \tag{3.35}$$

for all $h \in W_0^{1,p}(\Omega)$. In (3.35) we first choose $h = -\tilde{u}^- \in W_0^{1,p}(\Omega)$ and obtain $\tilde{u} \geq 0$ and $\tilde{u} \neq 0$. Then we choose $h = (\tilde{u} - u)^+ \in W_0^{1,p}(\Omega)$. This yields by applying (3.33) along with (3.29) and the fact that $u \in \mathcal{S}_+$

$$\begin{aligned} & \langle A_p(\tilde{u}), (\tilde{u} - u)^+ \rangle + \langle A_q(\tilde{u}), (\tilde{u} - u)^+ \rangle \\ &= \int_{\Omega} [\hat{c}_2 u^{\tau-1} - \hat{c}_1 u^{r-1}] (\tilde{u} - u)^+ dx \\ &\leq \int_{\Omega} [\vartheta(x) u^{\tau-1} + f(x, u)] (\tilde{u} - u)^+ dx \\ &= \langle A_p(u), (\tilde{u} - u)^+ \rangle + \langle A_q(u), (\tilde{u} - u)^+ \rangle. \end{aligned}$$

Hence, $\tilde{u} \leq u$. So we have proved that

$$\tilde{u} \in [0, u], \tilde{u} \neq 0. \tag{3.36}$$

From (3.36), (3.33), and (3.35), it follows that \tilde{u} is a positive solution of (3.30). Then $\tilde{u} = \bar{u} \in \text{int}(C_0^1(\bar{\Omega})_+)$ and so $\bar{u} \leq u$ for all $u \in \mathcal{S}_+$.

Similarly, we show that $v \leq \bar{v}$ for all $v \in \mathcal{S}_-$.

We have the following corollary.

Corollary 3.12 *Let hypotheses H_0 and H_1 be satisfied. Then*

$$\emptyset \neq \mathcal{S}_+ \subseteq \text{int}\left(C_0^1(\bar{\Omega})_+\right) \quad \text{and} \quad \emptyset \neq \mathcal{S}_- \subseteq -\text{int}\left(C_0^1(\bar{\Omega})_+\right).$$

Now we are ready to produce extremal constant sign solutions.

Proposition 3.13 *Let hypotheses H_0 and H_1 be satisfied. Then there exist solutions $u_* \in \mathcal{S}_+$ and $v_* \in \mathcal{S}_-$ such that*

$$u_* \leq u \text{ for all } u \in \mathcal{S}_+ \text{ and } v \leq v_* \text{ for all } v \in \mathcal{S}_-.$$

Proof From Papageorgiou et al. [19, Proposition 7], we know that \mathcal{S}_+ is downward directed. So, using Lemma 3.10 of Hu and Papageorgiou [8], we can find a decreasing sequence $\{u_n\}_{n \in \mathbb{N}}$ such that

$$\inf_{n \in \mathbb{N}} u_n = \inf \mathcal{S}_+.$$

Since $u_n \in \mathcal{S}_+$, we have

$$\langle A_p(u_n), h \rangle + \langle A_q(u_n), h \rangle = \int_{\Omega} \vartheta(x) u_n^{\tau-1} h \, dx + \int_{\Omega} f(x, u_n) h \, dx \tag{3.37}$$

for all $h \in W_0^{1,p}(\Omega)$. Evidently, the sequence $\{u_n\}_{n \in \mathbb{N}} \subseteq W_0^{1,p}(\Omega)$ is bounded. So, we may assume that

$$u_n \rightharpoonup u_* \text{ in } W_0^{1,p}(\Omega) \text{ and } u_n \rightarrow u_* \text{ in } L^r(\Omega). \tag{3.38}$$

Choosing $h = u_n - u$ in (3.37), passing to the limit as $n \rightarrow \infty$, and using the convergence properties in (3.38), we obtain

$$\limsup_{n \rightarrow \infty} \langle A_p(u_n), u_n - u \rangle \leq 0.$$

Then, by the (S_+) -property of A_p , we get

$$u_n \rightarrow u_* \text{ in } W_0^{1,p}(\Omega). \tag{3.39}$$

Passing to the limit in (3.37) and using (3.39), we have

$$\langle A_p(u_*), h \rangle + \langle A_q(u_*), h \rangle = \int_{\Omega} \vartheta(x) u_*^{\tau-1} h \, dx + \int_{\Omega} f(x, u_*) h \, dx$$

for all $h \in W_0^{1,p}(\Omega)$. From Proposition 3.11, we know that $\bar{u} \leq u_*$. Hence, $u_* \in \mathcal{S}_+$ and $u_* \leq u$ for all $u \in \mathcal{S}_+$.

Similarly, we produce $v_* \in \mathcal{S}_-$ such that $v \leq v_*$ for all $v \in \mathcal{S}_-$. Note that \mathcal{S}_- is upward directed.

Using the extremal constant sign solutions obtained in Proposition 3.13, we are going to prove the existence of a sign-changing solution. As explained earlier, we

focus on the order interval $[v_*, u_*]$ and look for solutions in $[v_*, u_*] \setminus \{0, u_*, v_*\}$. Such a solution turns out to be sign-changing.

Implementing the approach just described, let $u_* \in \mathcal{S}_+$ and $v_* \in \mathcal{S}_-$ be the extremal constant sign solutions from Proposition 3.13 and consider the truncation functions $k_1, k_2: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$k_1(x, s) = \begin{cases} \vartheta(x)|v_*(x)|^{\tau-2}v_*(x) & \text{if } s < v_*(x), \\ \vartheta(x)|s|^{\tau-2}s & \text{if } v_*(x) \leq s \leq u_*(x), \\ \vartheta(x)u_*(x)^{\tau-1} & \text{if } u_*(x) < s, \end{cases} \tag{3.40}$$

and

$$k_2(x, s) = \begin{cases} f(x, v_*(x)) & \text{if } s < v_*(x), \\ f(x, s) & \text{if } v_*(x) \leq s \leq u_*(x), \\ f(x, u_*(x)) & \text{if } u_*(x) < s. \end{cases} \tag{3.41}$$

It is clear that both are Carathéodory functions. We set

$$k(x, s) = k_1(x, s) + k_2(x, s). \tag{3.42}$$

Furthermore, we introduce the positive and negative truncations of $k(x, \cdot)$, namely the Carathéodory functions

$$k_{\pm}(x, s) = k_1(x, \pm s^{\pm}) + k_2(x, \pm s^{\pm}). \tag{3.43}$$

We set

$$\begin{aligned} K_1(x, s) &= \int_0^s k_1(x, t) dt, & K_2(x, s) &= \int_0^s k_2(x, t) dt, \\ K(x, s) &= K_1(x, s) + K_2(x, s), & K_{\pm}(x, s) &= \int_0^s k_{\pm}(x, t) dt \end{aligned}$$

and consider the C^1 -functionals $\zeta, \zeta_{\pm}: W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} \zeta(u) &= \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \int_{\Omega} K(x, u) dx \quad \text{for all } u \in W_0^{1,p}(\Omega) \tag{3.44} \\ \zeta_{\pm}(u) &= \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \int_{\Omega} K_{\pm}(x, u) dx \quad \text{for all } u \in W_0^{1,p}(\Omega), \end{aligned}$$

Applying (3.40), (3.41), (3.42), and (3.43), we check easily that

$$K_{\zeta} \subseteq [v_*, u_*] \cap C_0^1(\overline{\Omega}), \quad K_{\zeta_+} \subseteq [0, u_*] \cap C_0^1(\overline{\Omega}) \quad \text{and} \quad K_{\zeta_-} \subseteq [v_*, 0] \cap (-C_0^1(\overline{\Omega})).$$

Due to the extremality of u_* and v_* , we conclude that

$$K_\zeta \subseteq [v_*, u_*] \cap C_0^1(\bar{\Omega}), K_{\zeta_+} = \{0, u_*\} \text{ and } K_{\zeta_-} = \{0, v_*\}. \tag{3.45}$$

Proposition 3.14 *Let hypotheses H_0 and H_1 be satisfied. Then $u_* \in \mathcal{S}_+$ and $v_* \in \mathcal{S}_-$ are local minimizers of $\zeta : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$.*

Proof Because of (3.40), (3.41), and (3.43), it is clear that $\zeta_+ : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ is coercive and it is also sequentially weakly lower semicontinuous. Hence, we find $\tilde{u}_* \in W_0^{1,p}(\Omega)$ such that

$$\zeta_+(\tilde{u}_*) = \inf \left[\zeta_+(u) : u \in W_0^{1,p}(\Omega) \right] < 0 = \zeta_+(0), \tag{3.46}$$

since $\tau < q < p$, for $t \in (0, 1)$ small enough, we have by using H_1 (iv) and choosing $\varepsilon \in (0, \eta - \|\vartheta\|_\infty)$

$$\zeta_+(tu_*) \leq t^p \frac{\|\nabla u_*\|_p^p}{p} + t^q \frac{\|\nabla u_*\|_q^q}{q} + t^\tau \frac{1}{\tau} \left(\int_\Omega [\|\vartheta\|_\infty - (\eta - \varepsilon)] u_*^\tau dx \right) < 0.$$

Due to (3.46), we know that $\tilde{u}_* \in K_{\zeta_+}$ and so $\tilde{u}_* = u_*$, see (3.45). Let $\varrho > 0$ and

$$\overline{B}_\varrho^{C_0^1} = \left\{ u \in C_0^1(\bar{\Omega}) : \|u - u_*\|_{C_0^1(\bar{\Omega})} \leq \varrho \right\}.$$

Since $\zeta|_{C_0^1(\bar{\Omega})_+} = \zeta_+|_{C_0^1(\bar{\Omega})_+}$, we obtain for $u \in \overline{B}_\varrho^{C_0^1}$

$$\begin{aligned} \zeta(u) - \zeta(u_*) &= \zeta(u) - \zeta_+(u_*) \\ &\geq \zeta(u) - \zeta_+(u) \\ &= \int_\Omega [K_+(x, u) - K(x, u)] dx \\ &= \int_\Omega -K_1(x, -u^-) dx + \int_\Omega -K_2(x, -u^-) dx. \end{aligned} \tag{3.47}$$

We write as abbreviation

$$\begin{aligned} \{-u^- < v_*\} &:= \{x \in \Omega : -u^-(x) < v_*(x)\}, \\ \{v_* \leq -u^-\} &:= \{x \in \Omega : v_*(x) \leq -u^-(x)\}. \end{aligned}$$

Then, for the first integral on the right-hand side in (3.47), we have

$$\begin{aligned}
 & \int_{\Omega} -K_1(x, -u^-) \, dx \\
 &= \int_{\{-u^- < v_*\}} \left(-\frac{\vartheta(x)}{\tau} |v_*|^\tau - \vartheta(x) \left[|v_*|^{\tau-2} v_* (-u^- - v_*) \right] \right) \, dx \\
 &+ \int_{\{v_* \leq -u^-\}} \frac{-\vartheta(x)}{\tau} (u^-)^\tau \, dx \\
 &\geq \int_{\{v_* \leq -u^-\}} \frac{-\vartheta(x)}{\tau} (u^-)^\tau \, dx.
 \end{aligned} \tag{3.48}$$

From H_1 (iv), for given $\varepsilon > 0$, we can find $\hat{c}_{11} = \hat{c}_{11}(\varepsilon) > 0$ such that

$$F(x, s) \leq \frac{\varepsilon - \tilde{c}}{\beta} |s|^\beta + \hat{c}_{11} \left(\lambda |s|^{\hat{\mu}(\lambda)} + |s|^r \right) \tag{3.49}$$

for a. a. $x \in \Omega$ and for all $s \in \mathbb{R}$. Using (3.49), the second integral on the right-hand side in (3.47) can be estimated by (see also the proof of Proposition 3.4)

$$\begin{aligned}
 & \int_{\Omega} -K_2(x, -u^-) \, dx \\
 &= \int_{\{-u^- < v_*\}} -[F(x, v_*) + f(x, v_*)(-u^- - v_*)] \, dx \\
 &- \int_{\{v_* \leq -u^-\}} F(x, -u^-) \, dx \\
 &\geq \int_{\{-u^- < v_*\}} -[F(x, v_*) + f(x, v_*)(-u^- - v_*)] \, dx \\
 &- \int_{\{v_* \leq -u^-\}} \xi_\lambda (\|u^-\|_\infty) (u^-)^p \, dx.
 \end{aligned} \tag{3.50}$$

Combining (3.47), (3.48), (3.50) and applying hypotheses H_0 , we obtain

$$\begin{aligned}
 & \zeta(u) - \zeta(u_*) \\
 &\geq \int_{\{-u^- < v_*\}} -[F(x, v_*) + f(x, v_*)(-u^- - v_*)] \, dx \\
 &+ \int_{\{v_* \leq -u^-\}} \left(\frac{-\vartheta(x)}{\tau} (u^-)^\tau - \xi_\lambda (\|u^-\|_\infty) (u^-)^p \right) \, dx \\
 &\geq \int_{\{-u^- < v_*\}} -[F(x, v_*) + f(x, v_*)(-u^- - v_*)] \, dx \\
 &+ \int_{\{v_* \leq -u^-\}} \left(\frac{1}{\tau} c_0 (u^-)^\tau - \xi_\lambda (\|u^-\|_\infty) (u^-)^p \right) \, dx.
 \end{aligned} \tag{3.51}$$

Recall that $u_* \in C_0^1(\bar{\Omega})_+ \setminus \{0\}$ and $u \in \bar{B}_\varrho^{C_0^1}$. Hence, we have

$$\|u^-\|_\infty \rightarrow 0 \text{ as } \varrho \rightarrow 0^+.$$

Thus, $|\{-u^- \leq v_*\}|_N \rightarrow 0$ as $\varrho \rightarrow 0^+$ and $|\{v_* \leq -u^-\}|_N > 0$ for $\varrho > 0$ small enough and it is also decreasing in ϱ . Then, for λ small and for $\varrho > 0$ small enough, from (3.51), it follows that u_* is a local $C_0^1(\bar{\Omega})$ -minimizer of ζ and from Papageorgiou and Rădulescu [16], we deduce that u_* is a local $W_0^{1,p}(\Omega)$ -minimizer of ζ .

Similarly, working with ζ_- instead of ζ_+ , we can show the result for $v_* \in \mathcal{S}_-$.

Now we are ready to generate a sign-changing solution for problem (1.1).

Proposition 3.15 *Let hypotheses H_0 and H_1 be satisfied. Then problem (1.1) has a sign-changing solution $y_0 \in [v_*, u_*] \cap C_0^1(\bar{\Omega})$.*

Proof We assume that K_ζ is finite, otherwise on account of (3.45), (3.40), and (3.41), we would have infinity smooth sign-changing solutions. Moreover, we assume that $\zeta(v_*) \leq \zeta(u_*)$. The analysis is similar if the opposite inequality holds. From Proposition 3.14, we know that u_* is a local minimizer of ζ . Recall that the functional ζ is coercive. So, it satisfies the C-condition, see, for example, Papageorgiou et al. [21, p.369]. So, using Theorem 5.7.6 of Papageorgiou et al. [21], we can find $\rho \in (0, 1)$ small enough such that

$$\zeta(v_*) \leq \zeta(u_*) < \inf [\zeta(u) : \|u - u_*\| = \rho] =: m_\rho \text{ and } \|v_* - u_*\| > \rho.$$

Therefore, we can use the mountain pass theorem and find $y_0 \in W_0^{1,p}(\Omega)$ such that

$$y_0 \in K_\zeta \subseteq [v_*, u_*] \cap C_0^1(\bar{\Omega}), \tag{3.52}$$

see (3.45), and

$$\zeta(v_*) \leq \zeta(u_*) < m_\rho \leq \zeta(y_0). \tag{3.53}$$

From (3.53), we see that $y_0 \notin \{v_*, u_*\}$. Moreover, Theorem 6.5.8 of Papageorgiou et al. [21] implies that

$$C_1(\zeta, y_0) \neq 0. \tag{3.54}$$

On the other hand, the presence of the concave term and the C^1 -continuity of critical groups imply that

$$C_k(\zeta, 0) = 0 \text{ for all } k \in \mathbb{N}_0, \tag{3.55}$$

see Leonardi and Papageorgiou [11, Proposition 6] and Papageorgiou et al. [21, Proposition 6.3.4]. Comparing (3.54) and (3.55), we infer that $y_0 \neq 0$. Taking (3.52) into account, we conclude that y_0 is a smooth sign-changing solution of problem (1.1).

Summarizing this, we can state the following multiplicity theorem for problem (1.1).

Theorem 3.16 *Let hypotheses H_0 and H_1 be satisfied. Then problem (1.1) has at least three nontrivial smooth solutions*

$$u_0 \in \text{int} \left(C_0^1(\overline{\Omega})_+ \right), \quad v_0 \in -\text{int} \left(C_0^1(\overline{\Omega})_+ \right)$$

and

$$y_0 \in [v_0, u_0] \cap C_0^1(\overline{\Omega}) \text{ being sign-changing.}$$

4 Infinitely Many Nodal Solutions

In this section, under a local symmetry condition on $f(x, \cdot)$, we prove the existence of a whole sequence of nodal solutions converging to 0 in $C_0^1(\overline{\Omega})$.

The new conditions on the perturbation $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ are the following ones:

H_2 : $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function such that $f(x, \cdot)$ is odd for a. a. $x \in \Omega$ in $[-\gamma, \gamma]$ with $\gamma > 0$ and it satisfies the following assumptions:

- (i) there exist $r \in (p, p^*)$ and $0 \leq a(\cdot) \in L^\infty(\Omega)$ such that

$$|f(x, s)| \leq a(x) \left(1 + |s|^{r-1} \right)$$

for a. a. $x \in \Omega$ and for all $s \in \mathbb{R}$;

- (ii) if $F(x, s) = \int_0^s f(x, t) dt$, then

$$\lim_{s \rightarrow +\infty} \frac{F(x, s)}{s^p} = +\infty$$

uniformly for a. a. $x \in \Omega$ and there exists

$$\mu \in \left((r - p) \max \left\{ \frac{N}{p}, 1 \right\}, p^* \right)$$

such that

$$0 < \beta_0 \leq \liminf_{s \rightarrow +\infty} \frac{f(x, s)s - pF(x, s)}{s^\mu}$$

uniformly for a. a. $x \in \Omega$;

- (iii) there exist $\beta_1 \in L^\infty(\Omega)$ and $\beta_2 > 0$ such that

$$\hat{\lambda}_1(p) \leq \beta_1(x) \text{ for a. a. } x \in \Omega$$

with $\beta_1 \neq \hat{\lambda}_1(p)$ and

$$\beta_1(x) \leq \liminf_{s \rightarrow -\infty} \frac{f(x, s)}{|s|^{p-2}s} \leq \limsup_{s \rightarrow -\infty} \frac{f(x, s)}{|s|^{p-2}s} \leq \beta_2$$

uniformly for a. a. $x \in \Omega$.

(iv) there exists $\beta \in (1, \tau)$ such that

$$\lim_{s \rightarrow 0} \frac{f(x, s)}{|s|^{\beta-2}s} = 0$$

uniformly for a. a. $x \in \Omega$ and

$$\liminf_{s \rightarrow 0} \frac{f(x, s)}{|s|^{\tau-2}s} \geq \eta > \|\vartheta\|_\infty$$

uniformly for a. a. $x \in \Omega$ and for every $\lambda > 0$ there exists $\hat{\mu}(\lambda) \in (1, \beta)$ such that $\hat{\mu}(\lambda) \rightarrow \hat{\mu} \in (1, \beta)$ as $\lambda \rightarrow 0^+$ and

$$f(x, s)s \leq \hat{c} \left(\lambda |s|^{\hat{\mu}(\lambda)} + |s|^r \right) - \tilde{c} |s|^\beta$$

for a. a. $x \in \Omega$, for all $s \in \mathbb{R}$ with $\hat{c}, \tilde{c} > 0$.

Recall that the functional $\zeta : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ is given by

$$\zeta(u) = \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \int_\Omega K(x, u) \, dx \quad \text{for all } u \in W_0^{1,p}(\Omega),$$

see (3.44), where the difference is that, due to the local oddness of $f(x, \cdot)$, we truncate in (3.40), (3.41) above at $\text{int}(C_0^1(\overline{\Omega})_+) \ni \hat{\eta} < \min\{\gamma, u_*\}$ instead of u_* and below at $-\text{int}(C_0^1(\overline{\Omega})_+) \ni (-\hat{\eta}) > \max\{-\gamma, v_*\}$ instead of v_* . Let $V \subseteq W_0^{1,p}(\Omega) \cap L^\infty(\Omega)$ be a finite-dimensional subspace.

Proposition 4.1 *Let hypotheses H_0 and H_2 be satisfied. Then there exists $\rho_V > 0$ such that*

$$\sup[\zeta(u) : u \in V, \|u\| = \rho_V] < 0.$$

Proof On account of hypothesis H_2 (iv), for a given $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon) > 0$ such that

$$F(x, s) \geq \frac{1}{\tau} (\eta - \varepsilon) |s|^\tau \tag{4.1}$$

for a. a. $x \in \Omega$ and for all $|s| \leq \delta$.

Since V is finite dimensional, all norms are equivalent. Therefore, we can find $\rho_V > 0$ such that

$$u \in V \text{ and } \|u\| \leq \rho_V \text{ imply } |u(x)| \leq \delta \text{ for a. a. } x \in \Omega. \tag{4.2}$$

Applying (4.1) and (4.2), we have for $\|u\| \leq \rho_V$

$$\zeta(u) \leq \frac{1}{p} \|\nabla u\|^p + \frac{1}{q} \|\nabla u\|^q - \frac{1}{\tau} \int_{\Omega} (\eta - \varepsilon - \|\vartheta\|_{\infty}) |u|^{\tau} dx,$$

see the truncations in (3.40) and (3.41). Recalling that $\eta > \|\vartheta\|_{\infty}$, we choose $\varepsilon \in (0, \eta - \|\vartheta\|_{\infty})$. Then, using once more the fact that on V all norms are equivalent, we obtain

$$\zeta(u) \leq \frac{1}{p} \|\nabla u\|^p + \frac{1}{q} \|\nabla u\|^q - \hat{c}_1 \|u\|^{\tau}$$

for some $\hat{c}_1 > 0$.

Since $\tau < q < p$, choosing $\rho_V \in (0, 1)$ even smaller if necessary, we have

$$\sup \{ \zeta(u) : u \in V, \|u\| = \rho_V \} < 0.$$

Now we are ready for the new multiplicity theorem for problem (1.1) under H_2 .

Theorem 4.2 *Let hypotheses H_0 and H_2 be satisfied. Then problem (1.1) has a whole sequence of distinct nodal solutions $\{u_n\}_{n \in \mathbb{N}}$ such that $u_n \rightarrow 0$ in $C_0^1(\bar{\Omega})$.*

Proof Evidently, the functional $\zeta : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$ is even, $\zeta(0) = 0$ and it is bounded below and satisfies the C-condition being coercive due to (3.40) as well as (3.41). Then it satisfies the PS-condition as well, see Papageorgiou et al. [21, Proposition 5.1.14]. On account of Proposition 4.1, we can apply Theorem 1 of Kajikiya [9] and obtain a sequence $\{u_n\}_{n \in \mathbb{N}} \subseteq W_0^{1,p}(\Omega)$ such that

$$u_n \in K_{\zeta} \text{ for all } n \in \mathbb{N} \text{ and } u_n \rightarrow 0 \text{ in } W_0^{1,p}(\Omega).$$

Note that $u_n \in L^{\infty}(\Omega)$ (see, for example Ho et al. [7, Theorem 3.1]). Then, from the nonlinear regularity theory due to Lieberman [12, p. 320], there exist $\alpha \in (0, 1)$ and $M > 0$ such that

$$u_n \in C_0^{1,\alpha}(\bar{\Omega}) \text{ and } \|u_n\|_{C_0^{1,\alpha}(\bar{\Omega})} \leq M.$$

Using the compactness of $C_0^{1,\alpha}(\bar{\Omega})$ into $C_0^1(\bar{\Omega})$ gives

$$u_n \in C_0^1(\bar{\Omega}) \text{ for all } n \in \mathbb{N} \text{ and } u_n \rightarrow 0 \text{ in } C_0^1(\bar{\Omega}).$$

Since $\text{int}_{C_0^1(\bar{\Omega})} [v_*, u_*] \neq \emptyset$ (recall that $v_* \in -\text{int}(C_0^1(\bar{\Omega})_+)$, $u_* \in \text{int}(C_0^1(\bar{\Omega})_+)$), it follows that $\{u_n\}_{n \geq n_0} \subseteq [v_*, u_*]$ for some $n_0 \in \mathbb{N}$. These are nodal solutions of (1.1).

Remark 4.3 It will be interesting to extend the results of this paper to anisotropic equations. We believe that this is feasible. However, concerning possible extensions to double-phase problems with unbalanced growth, we doubt that this is possible due to the lack of a global regularity theory for such problems.

Acknowledgements The authors wish to thank the two anonymous referees for their comments and remarks which helped to improve the presentation of the paper. The first author was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - The Berlin Mathematics Research Center MATH+ and the Berlin Mathematical School (BMS) (EXC-2046/1, project ID: 390685689).

Funding Open Access funding enabled and organized by Projekt DEAL.

Declarations

Conflict of interest The authors have no conflict of interest to declare that is relevant to the content of this article.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Aizicovici, S., Papageorgiou, N.S., Staicu, V.: Degree theory for operators of monotone type and nonlinear elliptic equations with inequality constraints. *Mem. Am. Math. Soc.* **196**(915), 70 (2008)
2. Ambrosetti, A., Brezis, H., Cerami, G.: Combined effects of concave and convex nonlinearities in some elliptic problems. *J. Funct. Anal.* **122**(2), 519–543 (1994)
3. de Paiva, F.O., Massa, E.: Multiple solutions for some elliptic equations with a nonlinearity concave at the origin. *Nonlinear Anal.* **66**(12), 2940–2946 (2007)
4. Díaz, J.I., Saá, J.E.: Existence et unicité de solutions positives pour certaines équations elliptiques quasilineaires. *C. R. Acad. Sci. Paris Sér. I Math.* **305**(12), 521–524 (1987)
5. García Azorero, J.P., Peral Alonso, I., Manfredi, J.J.: Sobolev versus Hölder local minimizers and global multiplicity for some quasilinear elliptic equations. *Commun. Contemp. Math.* **2**(3), 385–404 (2000)
6. Ghoussoub, N.: *Duality and Perturbation Methods in Critical Point Theory*. Cambridge University Press, Cambridge (1993)
7. Ho, K., Kim, Y.-H., Winkert, P., Zhang, C.: The boundedness and Hölder continuity of solutions to elliptic equations involving variable exponents and critical growth. *J. Differ. Equ.* **313**, 503–532 (2022)
8. Hu, S., Papageorgiou, N.S.: *Handbook of Multivalued Analysis*, vol. I. Kluwer Academic Publishers, Dordrecht (1997)
9. Kajikiya, R.: A critical point theorem related to the symmetric mountain pass lemma and its applications to elliptic equations. *J. Funct. Anal.* **225**(2), 352–370 (2005)
10. Lê, A.: Eigenvalue problems for the p -Laplacian. *Nonlinear Anal.* **64**(5), 1057–1099 (2006)
11. Leonardi, S., Papageorgiou, N.S.: On a class of critical Robin problems. *Forum Math.* **32**(1), 95–109 (2020)
12. Lieberman, G.M.: The natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva for elliptic equations. *Commun. Partial Differ. Equ.* **16**(2–3), 311–361 (1991)

13. Marano, S.A., Mosconi, S.: Some recent results on the Dirichlet problem for (p, q) -Laplace equations. *Discret. Contin. Dyn. Syst. Ser. S* **11**(2), 279–291 (2018)
14. Motreanu, D., Motreanu, V.V., Papageorgiou, N.S.: *Topological and Variational Methods with Applications to Nonlinear Boundary Value Problems*. Springer, New York (2014)
15. Papageorgiou, N.S., Winkert, P.: Resonant $(p, 2)$ -equations with concave terms. *Appl. Anal.* **94**(2), 342–360 (2015)
16. Papageorgiou, N.S., Rădulescu, V.D.: Nonlinear nonhomogeneous Robin problems with superlinear reaction term. *Adv. Nonlinear Stud.* **16**(4), 737–764 (2016)
17. Papageorgiou, N.S., Winkert, P.: Asymmetric $(p, 2)$ -equations, superlinear at $+\infty$, resonant at $-\infty$. *Bull. Sci. Math.* **141**(5), 443–488 (2017)
18. Papageorgiou, N.S., Winkert, P.: *Applied Nonlinear Functional Analysis*. De Gruyter, Berlin (2018)
19. Papageorgiou, N.S., Rădulescu, V.D., Repovš, D.D.: Positive solutions for perturbations of the Robin eigenvalue problem plus an indefinite potential. *Discret. Contin. Dyn. Syst.* **37**(5), 2589–2618 (2017)
20. Papageorgiou, N.S., Rădulescu, V.D., Repovš, D.D.: Asymmetric Robin problems with indefinite potential and concave terms. *Adv. Nonlinear Stud.* **19**(1), 69–87 (2019)
21. Papageorgiou, N.S., Rădulescu, V.D., Repovš, D.D.: *Nonlinear Analysis-Theory and Methods*. Springer, Cham (2019)
22. Perera, K.: Multiplicity results for some elliptic problems with concave nonlinearities. *J. Differ. Equ.* **140**(1), 133–141 (1997)
23. Pucci, P., Serrin, J.: *The Maximum Principle*. Birkhäuser Verlag, Basel (2007)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.