



On a class of singular anisotropic (p, q) -equations

Nikolaos S. Papageorgiou¹ · Patrick Winkert²

Received: 22 September 2020 / Accepted: 23 March 2021
© The Author(s) 2021

Abstract

We consider a Dirichlet problem driven by the anisotropic (p, q) -Laplacian and with a reaction that has the competing effects of a singular term and of a parametric super-linear perturbation. Based on variational tools along with truncation and comparison techniques, we prove a bifurcation-type result describing the changes in the set of positive solutions as the parameter varies.

Keywords Anisotropic (p, q) -Laplacian · Singular term · Superlinear perturbation · Regularity theory · Maximum principle · Positive solutions

Mathematics Subject Classification 35J75

1 Introduction

Let $\Omega \subseteq \mathbb{R}^N$ be a bounded domain with a C^2 -boundary $\partial\Omega$. In this paper, we study the following anisotropic Dirichlet problem

$$\begin{aligned} -\Delta_{p(\cdot)}u - \Delta_{q(\cdot)}u &= u^{-\eta(x)} + \lambda f(x, u) \quad \text{in } \Omega \\ u|_{\partial\Omega} &= 0, \quad u > 0, \quad \lambda > 0. \end{aligned} \tag{P_\lambda}$$

✉ Patrick Winkert
winkert@math.tu-berlin.de
Nikolaos S. Papageorgiou
npapg@math.ntua.gr

¹ Department of Mathematics, National Technical University, Zografou Campus, Athens 15780, Greece

² Technische Universität Berlin, Institut für Mathematik, Straße des 17. Juni 136, 10623 Berlin, Germany

For $r \in E_1$, where E_1 is given by

$$E_1 = \left\{ r \in C(\overline{\Omega}) : 1 < \min_{x \in \overline{\Omega}} r(x) \right\},$$

we denote by $\Delta_{r(\cdot)}$ the anisotropic r -Laplacian (or $r(\cdot)$ -Laplacian) defined by

$$\Delta_{r(\cdot)} u = \operatorname{div} \left(|\nabla u|^{r(x)-2} \nabla u \right) \quad \text{for all } u \in W_0^{1,r(\cdot)}(\Omega).$$

The differential operator in problem (P_λ) is the sum of two such operators. In the reaction, the right-hand side of (P_λ) , we have the competing effects of two terms which are of different nature. One is the singular term $s \rightarrow s^{-\eta(x)}$ for $s > 0$ with $\eta \in C(\overline{\Omega})$ such that $0 < \eta(x) < 1$ for all $x \in \overline{\Omega}$. The other one is the parametric term $s \rightarrow \lambda f(x, s)$ with $\lambda > 0$ being the parameter and $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function, that is, $x \rightarrow f(x, s)$ is measurable for all $s \in \mathbb{R}$ and $s \rightarrow f(x, s)$ is continuous for a.a. $x \in \Omega$. We assume that $f(x, \cdot)$ exhibits $(p_+ - 1)$ -superlinear growth for a.a. $x \in \Omega$ as $s \rightarrow +\infty$ with $p_+ = \max_{x \in \overline{\Omega}} p(x)$. We are looking for positive solutions of problem (P_λ) and our aim is to determine how the set of positive solutions of (P_λ) changes as the parameter λ moves on the semiaxis $\mathring{\mathbb{R}}_+ = (0, +\infty)$.

The starting point of our work is the recent paper of Papageorgiou-Winkert [16] where the authors study a similar problem driven by the isotropic p -Laplacian. So, the differential operator in [16] is $(p - 1)$ -homogeneous and this property is exploited in their arguments. In contrast here, the differential operator is both nonhomogeneous and anisotropic.

Anisotropic problems with competition phenomena in the source were recently investigated by Papageorgiou-Rădulescu-Repovš [11]. They studied concave-convex problems driven by the $p(\cdot)$ -Laplacian plus an indefinite potential term. In their equation there is no singular term. In fact, the study of anisotropic singular problems is lagging behind. We are aware only the works of Byun-Ko [2] and Saoudi-Ghanmi [20] for Dirichlet as well as of Saoudi-Kratou-Alsadhan [21] for Neumann problems. All the aforementioned works deal with equations driven by the $p(\cdot)$ -Laplacian.

We mention that equations driven by the sum of two differential operators of different nature arise often in the mathematical models of physical processes. We mention the works of Bahrouni-Rădulescu-Repovš [1] (transonic flow problems), Cherfils-Il'yasov [3] (reaction diffusion systems) and Zhikov [26] (elasticity problems). Some recent regularity and multiplicity results can be found in the works of Ragusa-Tachikawa [19] and Papageorgiou-Zhang [17].

In this paper, under general conditions on the perturbation $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ which are less restrictive than all the previous cases in the literature, we prove the existence of a critical parameter $\lambda^* > 0$ such that

- for every $\lambda \in (0, \lambda^*)$, problem (P_λ) has at least two positive smooth solutions;
- for $\lambda = \lambda^*$, problem (P_λ) has at least one positive smooth solution;
- for every $\lambda > \lambda^*$, problem (P_λ) has no positive solutions.

2 Preliminaries and hypotheses

The study of anisotropic equations uses Lebesgue and Sobolev spaces with variable exponents. A comprehensive presentation of the theory of such spaces can be found in the book of Diening-Harjulehto-Hästö-Růžička [4].

Recall that $E_1 = \{r \in C(\overline{\Omega}) : 1 < \min_{x \in \overline{\Omega}} r(x)\}$. For any $r \in E_1$ we define

$$r_- = \min_{x \in \overline{\Omega}} r(x) \quad \text{and} \quad r_+ = \max_{x \in \overline{\Omega}} r(x).$$

Moreover, let $M(\Omega)$ be the space of all measurable functions $u : \Omega \rightarrow \mathbb{R}$. As usual, we identify two such functions when they differ only on a Lebesgue-null set. Then, given $r \in E_1$, the variable exponent Lebesgue space $L^{r(\cdot)}(\Omega)$ is defined as

$$L^{r(\cdot)}(\Omega) = \left\{ u \in M(\Omega) : \int_{\Omega} |u|^{r(x)} dx < \infty \right\}.$$

We equip this space with the so-called Luxemburg norm defined by

$$\|u\|_{r(\cdot)} = \inf \left\{ \lambda > 0 : \int_{\Omega} \left(\frac{|u|}{\lambda} \right)^{r(x)} dx \leq 1 \right\}.$$

Then $(L^{r(\cdot)}(\Omega), \|\cdot\|_{r(\cdot)})$ is a separable and reflexive Banach space, in fact it is uniformly convex. Let $r' \in E_1$ be the conjugate variable exponent to r , that is,

$$\frac{1}{r(x)} + \frac{1}{r'(x)} = 1 \quad \text{for all } x \in \Omega.$$

We know that $L^{r(\cdot)}(\Omega)^* = L^{r'(\cdot)}(\Omega)$ and the following Hölder type inequality holds

$$\int_{\Omega} |uv| dx \leq \left[\frac{1}{r_-} + \frac{1}{r'_-} \right] \|u\|_{r(\cdot)} \|v\|_{r'(\cdot)}$$

for all $u \in L^{r(\cdot)}(\Omega)$ and for all $v \in L^{r'(\cdot)}(\Omega)$.

If $r_1, r_2 \in E_1$ and $r_1(x) \leq r_2(x)$ for all $x \in \overline{\Omega}$, then we have that

$$L^{r_2(\cdot)}(\Omega) \hookrightarrow L^{r_1(\cdot)}(\Omega) \quad \text{continuously.}$$

The corresponding variable exponent Sobolev spaces can be defined in a natural way using the variable exponent Lebesgue spaces. So, if $r \in E_1$, then the variable exponent Sobolev space $W^{1,r(\cdot)}(\Omega)$ is defined by

$$W^{1,r(\cdot)}(\Omega) = \left\{ u \in L^{r(\cdot)}(\Omega) : |\nabla u| \in L^{r(\cdot)}(\Omega) \right\}.$$

Here the gradient ∇u is understood in the weak sense. We equip $W^{1,r(\cdot)}(\Omega)$ with the following norm

$$\|u\|_{1,r(\cdot)} = \|u\|_{r(\cdot)} + \|\nabla u\|_{r(\cdot)} \quad \text{for all } u \in W^{1,r(\cdot)}(\Omega).$$

In what follows we write $\|\nabla u\|_{r(\cdot)} = \|\nabla u\|_{r(\cdot)}$. Suppose that $r \in E_1$ is Lipschitz continuous, that is, $r_1 \in E_1 \cap C^{0,1}(\overline{\Omega})$. We define

$$W_0^{1,r(\cdot)}(\Omega) = \overline{C_c^\infty(\Omega)}^{\|\cdot\|_{1,r(\cdot)}}.$$

The spaces $W^{1,r(\cdot)}(\Omega)$ and $W_0^{1,r(\cdot)}(\Omega)$ are both separable and reflexive, in fact uniformly convex Banach spaces. On the space $W_0^{1,r(\cdot)}(\Omega)$ we have the Poincaré inequality, namely there exists $c_0 > 0$ such that

$$\|u\|_{r(\cdot)} \leq c_0 \|\nabla u\|_{r(\cdot)} \quad \text{for all } u \in W_0^{1,r(\cdot)}(\Omega).$$

Therefore, we can consider on $W_0^{1,r(\cdot)}(\Omega)$ the equivalent norm

$$\|u\|_{1,r(\cdot)} = \|\nabla u\|_{r(\cdot)} \quad \text{for all } u \in W_0^{1,r(\cdot)}(\Omega).$$

For $r \in E_1$ we introduce the critical Sobolev variable exponent r^* defined by

$$r^*(x) = \begin{cases} \frac{Nr(x)}{N-r(x)} & \text{if } r(x) < N, \\ +\infty & \text{if } N \leq r(x), \end{cases} \quad \text{for all } x \in \overline{\Omega}.$$

Suppose that $r \in E_1 \cap C^{0,1}(\overline{\Omega})$, $q \in E_1$, $q_+ < N$ and $1 < q(x) \leq r^*(x)$ for all $x \in \overline{\Omega}$. Then we have

$$W_0^{1,r(\cdot)}(\Omega) \hookrightarrow L^{q(\cdot)}(\Omega) \quad \text{continuously.}$$

Similarly, if $1 < q(x) < r^*(x)$ for all $x \in \overline{\Omega}$, we have

$$W_0^{1,r(\cdot)}(\Omega) \hookrightarrow L^{q(\cdot)}(\Omega) \quad \text{compactly.}$$

In the study of the variable exponent spaces, the modular function is important, that is, for $r \in E_1$,

$$\varrho_{r(\cdot)}(u) = \int_{\Omega} |u|^{r(x)} dx \quad \text{for all } u \in L^{r(\cdot)}(\Omega).$$

As before we write $\varrho_{r(\cdot)}(\nabla u) = \varrho_{r(\cdot)}(|\nabla u|)$. The importance of this function comes from the fact that it is closely related to the norm of the space. This is evident in the next proposition.

Proposition 2.1 *If $r \in E_1$, then we have the following assertions:*

$$(a) \quad \|u\|_{r(\cdot)} = \lambda \iff \varrho_{r(\cdot)}\left(\frac{u}{\lambda}\right) = 1 \text{ for all } u \in L^{r(\cdot)}(\Omega) \text{ with } u \neq 0;$$

- (b) $\|u\|_{r(\cdot)} < 1$ (resp. $= 1, > 1$) $\iff \varrho_{r(\cdot)}(u) < 1$ (resp. $= 1, > 1$);
- (c) $\|u\|_{r(\cdot)} < 1 \implies \|u\|_{r(\cdot)}^{r_+} \leq \varrho_{r(\cdot)}(u) \leq \|u\|_{r(\cdot)}^{r_-}$;
- (d) $\|u\|_{r(\cdot)} > 1 \implies \|u\|_{r(\cdot)}^{r_-} \leq \varrho_{r(\cdot)}(u) \leq \|u\|_{r(\cdot)}^{r_+}$;
- (e) $\|u_n\|_{r(\cdot)} \rightarrow 0 \iff \varrho_{r(\cdot)}(u_n) \rightarrow 0$;
- (f) $\|u_n\|_{r(\cdot)} \rightarrow +\infty \iff \varrho_{r(\cdot)}(u_n) \rightarrow +\infty$.

We know that for $r \in E_1 \cap C^{0,1}(\overline{\Omega})$, we have

$$W_0^{1,r(\cdot)}(\Omega)^* = W^{-1,r'(\cdot)}(\Omega).$$

Then we can introduce the nonlinear map $A_{r(\cdot)} : W_0^{1,r(\cdot)}(\Omega) \rightarrow W^{-1,r'(\cdot)}(\Omega)$ defined by

$$\langle A_{r(\cdot)}(u), h \rangle = \int_{\Omega} |\nabla u|^{r(x)-2} \nabla u \cdot \nabla h \, dx \quad \text{for all } u, h \in W_0^{1,r(\cdot)}(\Omega).$$

This map has the following properties, see, for example Gasiński-Papageorgiou [7, Proposition 2.5] and Rădulescu-Repovš [18, p. 40].

Proposition 2.2 *The operator $A_{r(\cdot)} : W_0^{1,r(\cdot)}(\Omega) \rightarrow W^{-1,r'(\cdot)}(\Omega)$ is bounded (so it maps bounded sets to bounded sets), continuous, strictly monotone (which implies it is also maximal monotone) and of type S_+ , that is,*

$$u_n \xrightarrow{w} u \text{ in } W_0^{1,r(\cdot)}(\Omega) \text{ and } \limsup_{n \rightarrow \infty} \langle A_{r(\cdot)}(u_n), u_n - u \rangle \leq 0$$

imply $u_n \rightarrow u$ in $W_0^{1,r(\cdot)}(\Omega)$.

Another space that we will use as a result of the anisotropic regularity theory is the Banach space

$$C_0^1(\overline{\Omega}) = \left\{ u \in C^1(\overline{\Omega}) : u|_{\partial\Omega} = 0 \right\}.$$

This is an ordered Banach space with positive (order) cone

$$C_0^1(\overline{\Omega})_+ = \left\{ u \in C_0^1(\overline{\Omega}) : u(x) \geq 0 \text{ for all } x \in \overline{\Omega} \right\}.$$

This cone has a nonempty interior given by

$$\text{int} \left(C_0^1(\overline{\Omega})_+ \right) = \left\{ u \in C_0^1(\overline{\Omega})_+ : u(x) > 0 \text{ for all } x \in \Omega, \frac{\partial u}{\partial n} \Big|_{\partial\Omega} < 0 \right\},$$

where $\frac{\partial u}{\partial n} = \nabla u \cdot n$ with n being the outward unit normal on $\partial\Omega$.

Let $h_1, h_2 \in M(\Omega)$. We write $h_1 \leq h_2$ if and only if $0 < c_K \leq h_2(x) - h_1(x)$ for a. a. $x \in K$ and for all compact sets $K \subseteq \Omega$. It is clear that if $h_1, h_2 \in C(\Omega)$ and

$h_1(x) < h_2(x)$ for all $x \in \Omega$, then $h_1 \preceq h_2$. From Papageorgiou-Rădulescu-Repovš [11, Proposition 2.4] and Papageorgiou-Rădulescu-Repovš [13, Propositions 6 and 7], we have the following comparison principles. In what follows, let $p, q \in E_1 \cap C^{0,1}(\overline{\Omega})$ with $q(x) < p(x)$ for all $x \in \overline{\Omega}$ and $\eta \in C(\overline{\Omega})$ with $0 < \eta(x) < 1$ for all $x \in \overline{\Omega}$.

Proposition 2.3

(a) If $\hat{\xi} \in L^\infty(\Omega)$, $\hat{\xi}(x) \geq 0$ for a. a. $x \in \Omega$, $h_1, h_2 \in L^\infty(\Omega)$, $h_1 \preceq h_2$, $u \in C_0^1(\overline{\Omega})_+$, $u > 0$ for all $x \in \Omega$, $v \in \text{int}(C_0^1(\overline{\Omega})_+)$ and

$$\begin{aligned} -\Delta_{p(\cdot)}u - \Delta_{q(\cdot)}u + \hat{\xi}(x)u^{p(x)-1} - u^{-\eta(x)} &= h_1(x) \text{ in } \Omega, \\ -\Delta_{p(\cdot)}v - \Delta_{q(\cdot)}v + \hat{\xi}(x)v^{p(x)-1} - v^{-\eta(x)} &= h_2(x) \text{ in } \Omega, \end{aligned}$$

then $v - u \in \text{int}(C_0^1(\overline{\Omega})_+)$.

(b) If $\hat{\xi} \in L^\infty(\Omega)$, $\hat{\xi} \geq 0$ for a. a. $x \in \Omega$, $h_1, h_2 \in L^\infty(\Omega)$, $0 < \hat{c} \leq h_2(x) - h_1(x)$ for a. a. $x \in \Omega$, $u, v \in C^1(\overline{\Omega}) \setminus \{0\}$, $u(x) \leq v(x)$ for all $x \in \Omega$, $v \in \text{int}(C_0^1(\overline{\Omega})_+)$ and

$$\begin{aligned} -\Delta_{p(\cdot)}u - \Delta_{q(\cdot)}u + \hat{\xi}(x)u^{p(x)-1} - u^{-\eta(x)} &= h_1(x) \text{ in } \Omega, \\ -\Delta_{p(\cdot)}v - \Delta_{q(\cdot)}v + \hat{\xi}(x)v^{p(x)-1} - v^{-\eta(x)} &= h_2(x) \text{ in } \Omega, \end{aligned}$$

then $u(x) < v(x)$ for all $x \in \Omega$.

Remark 2.4 Note that in part (a) of Proposition 2.3 we have by the weak comparison principle that $u \leq v$, see Tolksdorf [24].

If $u, v \in W_0^{1,p(\cdot)}(\Omega)$ with $u \leq v$, then we define

$$\begin{aligned} [u, v] &= \left\{ y \in W_0^{1,p(\cdot)}(\Omega) : u(x) \leq y(x) \leq v(x) \text{ for a. a. } x \in \Omega \right\}, \\ [u] &= \left\{ y \in W_0^{1,p(\cdot)}(\Omega) : u(x) \leq y(x) \text{ for a. a. } x \in \Omega \right\}. \end{aligned}$$

In what follows we will denote by $\|\cdot\|$ the norm of the Sobolev space $W_0^{1,p(\cdot)}(\Omega)$. By the Poincaré inequality we have

$$\|u\| = \|\nabla u\|_{p(\cdot)} \quad \text{for all } u \in W_0^{1,p(\cdot)}(\Omega).$$

Suppose that X is a Banach space and let $\varphi \in C^1(X)$. We denote the critical set of φ by

$$K_\varphi = \{u \in X : \varphi'(u) = 0\}.$$

Moreover, we say that φ satisfies the ‘‘Cerami condition’’, C-condition for short, if every sequence $\{u_n\}_{n \in \mathbb{N}} \subseteq X$ such that $\{\varphi(u_n)\}_{n \in \mathbb{N}} \subseteq \mathbb{R}$ is bounded and

$$(1 + \|u_n\|_X) \varphi'(u_n) \rightarrow 0 \text{ in } X^* \text{ as } n \rightarrow \infty,$$

admits a strongly convergent subsequence. This is a compactness-type condition on the functional φ which compensates for the fact that the ambient space X need not be locally compact being in general infinite dimensional. Applying this condition, one can prove a deformation theorem from which the minimax theorems for the critical values of φ follow. We refer to Papageorgiou-Rădulescu-Repovš [12, Chapter 5] and Struwe [22, Chapter II].

Given $s \in (1, +\infty)$ we denote by $s' \in (1, +\infty)$ the conjugate exponent defined by

$$\frac{1}{s} + \frac{1}{s'} = 1.$$

Furthermore, if $f: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a measurable function, then we denote by N_f the Nemytskii (also called superposition) operator corresponding to f , that is,

$$N_f(u)(\cdot) = f(\cdot, u(\cdot)) \text{ for all } u \in M(\Omega).$$

Note that $x \rightarrow f(x, u(x))$ is measurable. We know that if $f: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function, then $f(\cdot, \cdot)$ is jointly measurable, see Papageorgiou-Winkert [15, p. 106].

Now we are in the position to introduce our hypotheses on the data of problem (P_λ) .

H_0 : $p, q \in E_1 \cap C^{0,1}(\overline{\Omega})$, $\eta \in C(\overline{\Omega})$, $q(x) < p(x)$, $0 < \eta(x) < 1$ for all $x \in \overline{\Omega}$, $p_- < N$.

H_1 : $f: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function such that $f(x, 0) = 0$ for a. a. $x \in \Omega$ and

(i) there exists $a \in L^\infty(\Omega)$ such that

$$0 \leq f(x, s) \leq a(x) [1 + s^{r-1}]$$

for a. a. $x \in \Omega$, for all $s \geq 0$ and with $p_+ < r < p_-^*$, where

$$p_-^* = \frac{Np_-}{N - p_-};$$

(ii) if $F(x, s) = \int_0^s f(x, t) dt$, then

$$\lim_{s \rightarrow +\infty} \frac{F(x, s)}{s^{p_+}} = +\infty \text{ uniformly for a. a. } x \in \Omega;$$

(iii) there exists a function $\tau \in C(\overline{\Omega})$ such that

$$\tau(x) \in \left((r - p_-) \frac{N}{p_-}, p^*(x) \right) \quad \text{for all } x \in \overline{\Omega}$$

and

$$0 < \gamma_0 \leq \liminf_{s \rightarrow +\infty} \frac{f(x, s)s - p_+ F(x, s)}{s^{\tau(x)}} \quad \text{uniformly for a. a. } x \in \Omega;$$

(iv) for every $\rho > 0$ there exists $\hat{\xi}_\rho > 0$ such that the function

$$s \rightarrow f(x, s) + \hat{\xi}_\rho s^{p(x)-1}$$

is nondecreasing on $[0, \rho]$ for a. a. $x \in \Omega$.

Remark 2.5 Since we are interested in positive solutions and all the hypotheses above concern the positive semiaxis $\mathbb{R}_+ = [0, +\infty)$, we may assume without any loss of generality that $f(x, s) = 0$ for a. a. $x \in \Omega$ and for all $s \leq 0$. Hypotheses H_1 (ii), (iii) imply that $f(x, \cdot)$ is $(p_+ - 1)$ -superlinear for a. a. $x \in \Omega$. However, this superlinearity condition on $f(x, \cdot)$ is not formulated by using the Ambrosetti-Rabinowitz condition which is common in the literature when dealing with superlinear problems, see Byun-Ko [2], Saoudi-Ghanmi [20] and Saoudi-Kratou-Alsadhan [21]. Here, instead of the Ambrosetti-Rabinowitz condition, we employ hypothesis H_1 (iii) which is less restrictive and incorporates in our framework nonlinearities with “slower” growth near $+\infty$. For example, consider the functions

$$f_1(x, s) = (s + 1)^{p_+ - 1} \ln(s + 1) + s^{r_1(x) - 1} \quad \text{for all } s \geq 0$$

with $r_1 \in E_1, r_1(x) \leq p(x)$ for all $x \in \overline{\Omega}$ and

$$f_2(x, s) = \begin{cases} s^{\mu(x) - 1} & \text{if } 0 \leq s \leq 1, \\ s^{p_+ - 1} \ln(s) + s^{r_2(x) - 1} & \text{if } 1 < s \end{cases}$$

with $\mu, r_2 \in E_1$ and $r_2(x) \leq p(x)$ for all $x \in \overline{\Omega}$. These functions satisfy hypotheses H_1 , but fail to satisfy the Ambrosetti-Rabinowitz condition, see, for example, Gasiński-Papageorgiou [7].

The difficulty that we encounter when we study a singular problem is that the energy (Euler) functional of the problem is not C^1 because of the presence of the singular term. Hence, we cannot use the results of critical point theory. We need to find a way to bypass the singularity and deal with C^1 -functionals. In the next section, we examine a purely singular problem and the solution of this problem will help us in bypassing the singularity.

3 An auxiliary purely singular problem

In this section we deal with the following purely singular anisotropic (p, q) -equation

$$-\Delta_{p(\cdot)}u - \Delta_{q(\cdot)}u = u^{-\eta(x)} \quad \text{in } \Omega, \quad u|_{\partial\Omega} = 0, \quad u > 0. \tag{3.1}$$

Proposition 3.1 *If hypotheses H_0 hold, then problem (3.1) admits a unique position solution $\bar{u} \in \text{int}(C_0^1(\bar{\Omega})_+)$.*

Proof Let $g \in L^{p(\cdot)}(\Omega)$ and let $0 < \varepsilon \leq 1$. We consider the following Dirichlet problem

$$-\Delta_{p(\cdot)}u - \Delta_{q(\cdot)}u = [|g(x)| + \varepsilon]^{-\eta(x)} \quad \text{in } \Omega, \quad u|_{\partial\Omega} = 0, \quad u > 0.$$

Let $V : W_0^{1,p(\cdot)}(\Omega) \rightarrow W_0^{1,p(\cdot)}(\Omega)^* = W^{-1,p'(\cdot)}(\Omega)$ be the operator defined by

$$V(u) = A_{p(\cdot)}(u) + A_{q(\cdot)}(u) \quad \text{for all } u \in W_0^{1,p(\cdot)}(\Omega).$$

This map is continuous and strictly monotone, see Proposition 2.2, hence maximal monotone as well. It is also coercive, see Proposition 2.1. Therefore, it is surjective, see Papageorgiou-Rădulescu-Repovš [12, p. 135]. Since $[|g(\cdot)| + \varepsilon]^{-\eta(\cdot)} \in L^\infty(\Omega)$, there exists $u_\varepsilon \in W_0^{1,p(\cdot)}(\Omega)$, $u_\varepsilon \geq 0$, $u_\varepsilon \neq 0$ such that

$$V(u_\varepsilon) = [|g| + \varepsilon]^{-\eta(\cdot)}.$$

The strict monotonicity of V implies the uniqueness of u_ε . Thus, we can define the map $\beta : L^{p(\cdot)}(\Omega) \rightarrow L^{p(\cdot)}(\Omega)$ by setting

$$\beta(g) = u_\varepsilon.$$

Recall that $W_0^{1,p(\cdot)}(\Omega) \hookrightarrow L^{p(\cdot)}(\Omega)$ is compactly embedded. We claim that the map β is continuous. So, let $g_n \rightarrow g$ in $L^{p(\cdot)}(\Omega)$ and let $u_\varepsilon^n = \beta(g_n)$ with $n \in \mathbb{N}$. We have

$$\langle A_{p(\cdot)}(u_\varepsilon^n), h \rangle + \langle A_{q(\cdot)}(u_\varepsilon^n), h \rangle = \int_\Omega \frac{h}{[|g_n| + \varepsilon]^{\eta(x)}} dx \tag{3.2}$$

for all $h \in W_0^{1,p(\cdot)}(\Omega)$ and for all $n \in \mathbb{N}$.

We choose $h = u_\varepsilon^n \in W_0^{1,p(\cdot)}(\Omega)$ in (3.2) and obtain

$$\varrho_{p(\cdot)}(\nabla u_\varepsilon^n) + \varrho_{q(\cdot)}(\nabla u_\varepsilon^n) \leq \int_\Omega \frac{u_\varepsilon^n}{\varepsilon^{\eta_+}} dx,$$

which by Proposition 2.1 implies that

$$\{u_\varepsilon^n\}_{n \in \mathbb{N}} \subseteq W_0^{1,p(\cdot)}(\Omega) \text{ is bounded.}$$

So, we may assume that

$$u_\varepsilon^n \xrightarrow{w} \tilde{u}_\varepsilon \text{ in } W_0^{1,p(\cdot)}(\Omega) \text{ and } u_\varepsilon^n \rightarrow \tilde{u}_\varepsilon \text{ in } L^{p(\cdot)}(\Omega). \tag{3.3}$$

Now we choose $h = u_\varepsilon^n - \tilde{u}_\varepsilon \in W_0^{1,p(\cdot)}(\Omega)$ in (3.2), pass to the limit as $n \rightarrow \infty$ and apply (3.3) which results in

$$\lim_{n \rightarrow \infty} [\langle A_{p(\cdot)}(u_\varepsilon^n), u_\varepsilon^n - \tilde{u}_\varepsilon \rangle + \langle A_{q(\cdot)}(u_\varepsilon^n), u_\varepsilon^n - \tilde{u}_\varepsilon \rangle] = 0.$$

Since $A_{q(\cdot)}(\cdot)$ is monotone, we have

$$\limsup_{n \rightarrow \infty} [\langle A_{p(\cdot)}(u_\varepsilon^n), u_\varepsilon^n - \tilde{u}_\varepsilon \rangle + \langle A_{q(\cdot)}(\tilde{u}_\varepsilon), u_\varepsilon^n - \tilde{u}_\varepsilon \rangle] \leq 0.$$

Applying (3.3) gives

$$\limsup_{n \rightarrow \infty} \langle A_{p(\cdot)}(u_\varepsilon^n), u_\varepsilon^n - \tilde{u}_\varepsilon \rangle \leq 0$$

and so, by Proposition 2.2,

$$u_\varepsilon^n \rightarrow \tilde{u}_\varepsilon \text{ in } W_0^{1,p(\cdot)}(\Omega). \tag{3.4}$$

Passing to the limit in (3.2) as $n \rightarrow \infty$ and using (3.4) yields

$$\langle A_{p(\cdot)}(\tilde{u}_\varepsilon), h \rangle + \langle A_{q(\cdot)}(\tilde{u}_\varepsilon), h \rangle = \int_\Omega \frac{h}{[|g| + \varepsilon]^{\eta(x)}} dx$$

for all $h \in W_0^{1,p(\cdot)}(\Omega)$. Hence, $\tilde{u}_\varepsilon = \beta(g)$.

So, for the original sequence, we have

$$u_\varepsilon^n = \beta(g_n) \rightarrow \beta(g) = \tilde{u}_\varepsilon,$$

which shows that β is continuous.

From the argument above and recalling that $W_0^{1,p(\cdot)}(\Omega) \hookrightarrow L^{p(\cdot)}(\Omega)$ compactly, we see that $\beta(L^{p(\cdot)}(\Omega)) \subseteq L^{p(\cdot)}(\Omega)$ is compact. So, by the Schauder-Tychonov fixed point theorem, see Papageorgiou-Rădulescu-Repovš [12, p. 298] we can find $\hat{u}_\varepsilon \in W_0^{1,p(\cdot)}(\Omega)$ such that $\beta(\hat{u}_\varepsilon) = \hat{u}_\varepsilon$.

From Fan-Zhao [5], see also Gasiński-Papageorgiou [7] and Marino-Winkert [10], we have that $\hat{u}_\varepsilon \in L^\infty(\Omega)$. Then, from Tan-Fang [23, Corollary 3.1], we have $\hat{u}_\varepsilon \in C_0^1(\bar{\Omega}) \setminus \{0\}$. Finally, the anisotropic maximum principle of Zhang [25], see also Papageorgiou-Vetro-Vetro [14], implies that $\hat{u}_\varepsilon \in \text{int}(C_0^1(\bar{\Omega})_+)$.

Claim: If $0 < \varepsilon' \leq \varepsilon$, then $\hat{u}_\varepsilon \leq \hat{u}_{\varepsilon'}$. We have

$$-\Delta_{p(\cdot)}\hat{u}_{\varepsilon'} - \Delta_{q(\cdot)}\hat{u}_{\varepsilon'} = \frac{1}{[\hat{u}_{\varepsilon'} + \varepsilon']^{\eta(x)}} \geq \frac{1}{[\hat{u}_{\varepsilon'} + \varepsilon]^{\eta(x)}} \text{ in } \Omega. \tag{3.5}$$

We introduce the Carathéodory function $k_\varepsilon : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$k_\varepsilon(x, s) = \begin{cases} \frac{1}{[s^+ + \varepsilon]^{\eta(x)}} & \text{if } s \leq \hat{u}_{\varepsilon'}(x), \\ \frac{1}{[\hat{u}_{\varepsilon'}(x) + \varepsilon]^{\eta(x)}} & \text{if } \hat{u}_{\varepsilon'}(x) < s. \end{cases} \tag{3.6}$$

We set $K_\varepsilon(x, s) = \int_0^s k_\varepsilon(x, t) dt$ and consider the C^1 -functional $J_\varepsilon : W_0^{1,p(\cdot)}(\Omega) \rightarrow \mathbb{R}$ defined by

$$J_\varepsilon(u) = \int_\Omega \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_\Omega \frac{1}{q(x)} |\nabla u|^{q(x)} dx - \int_\Omega K_\varepsilon(x, u) dx$$

for all $u \in W_0^{1,p(\cdot)}(\Omega)$. From (3.6) it is clear that $J_\varepsilon : W_0^{1,p(\cdot)}(\Omega) \rightarrow \mathbb{R}$ is coercive and by the compact embedding $W_0^{1,p(\cdot)}(\Omega) \hookrightarrow L^r(\Omega)$ we know that it is also sequentially weakly lower semicontinuous. Therefore, by the Weierstraß-Tonelli theorem, there exists $\hat{u}_\varepsilon^* \in W_0^{1,p(\cdot)}(\Omega)$ such that

$$J_\varepsilon(\hat{u}_\varepsilon^*) = \min \left[J_\varepsilon(u) : u \in W_0^{1,p(\cdot)}(\Omega) \right]. \tag{3.7}$$

Let $u \in \text{int}(C_0^1(\overline{\Omega})_+)$ and choose $t \in (0, 1)$ small enough so that $tu \leq \hat{u}_{\varepsilon'}$, recall that $\hat{u}_{\varepsilon'} \in \text{int}(C_0^1(\overline{\Omega})_+)$ and use Proposition 4.1.22 of Papageorgiou-Rădulescu-Repovš [12]. Then, by (3.6), we obtain

$$\begin{aligned} J_\varepsilon(tu) &\leq \frac{t^{q-}}{q-} [\varrho_{p(\cdot)}(\nabla u) + \varrho_{q(\cdot)}(\nabla u)] - \int_\Omega \frac{1}{1 - \eta(x)} (tu)^{1-\eta(x)} dx \\ &\leq c_1 t^{q-} - c_2 t^{1-\eta-} \end{aligned}$$

for some $c_1 = c_1(u) > 0$, $c_2 = c_2(u) > 0$ and $t \in (0, 1)$. Choosing $t \in (0, 1)$ even smaller if necessary, we see that

$$J_\varepsilon(tu) < 0,$$

since $1 - \eta- < 1 < q-$. Then, by (3.7), because $\hat{u}_\varepsilon^* \in W_0^{1,p(\cdot)}(\Omega)$ is the global minimizer of J_ε , we conclude that

$$J_\varepsilon(\hat{u}_\varepsilon^*) < 0 = J_\varepsilon(0)$$

and so $\hat{u}_\varepsilon^* \neq 0$.

From (3.7) we have $J'_\varepsilon(\hat{u}_\varepsilon^*) = 0$ which means

$$\langle A_{p(\cdot)}(\hat{u}_\varepsilon^*), h \rangle + \langle A_{q(\cdot)}(\hat{u}_\varepsilon^*), h \rangle = \int_\Omega k_\varepsilon(x, \hat{u}_\varepsilon^*) dx \tag{3.8}$$

for all $h \in W_0^{1,p(\cdot)}(\Omega)$. Testing (3.8) with $h = -(\hat{u}_\varepsilon^*)^- \in W_0^{1,p(\cdot)}(\Omega)$ we obtain

$$\varrho_{p(\cdot)}(\nabla(\hat{u}_\varepsilon^*)^-) \leq 0,$$

because of (3.6) which by Proposition 2.1 implies that

$$\hat{u}_\varepsilon^* \geq 0 \quad \text{and} \quad \hat{u}_\varepsilon^* \neq 0.$$

Now we choose $h = (\hat{u}_\varepsilon^* - \hat{u}_{\varepsilon'})^+ \in W_0^{1,p(\cdot)}(\Omega)$ in (3.8). Applying (3.6) and (3.5) gives

$$\begin{aligned} & \left\langle A_{p(\cdot)}(\hat{u}_\varepsilon^*), (\hat{u}_\varepsilon^* - \hat{u}_{\varepsilon'})^+ \right\rangle + \left\langle A_{q(\cdot)}(\hat{u}_\varepsilon^*), (\hat{u}_\varepsilon^* - \hat{u}_{\varepsilon'})^+ \right\rangle \\ &= \int_\Omega \frac{1}{[\hat{u}_{\varepsilon'} + \varepsilon]^{\eta(x)}} (\hat{u}_\varepsilon^* - \hat{u}_{\varepsilon'})^+ dx \\ &\leq \left\langle A_{p(\cdot)}(\hat{u}_{\varepsilon'}), (\hat{u}_\varepsilon^* - \hat{u}_{\varepsilon'})^+ \right\rangle + \left\langle A_{q(\cdot)}(\hat{u}_{\varepsilon'}), (\hat{u}_\varepsilon^* - \hat{u}_{\varepsilon'})^+ \right\rangle. \end{aligned}$$

Hence, $\hat{u}_\varepsilon^* \leq \hat{u}_{\varepsilon'}$. So we have proved that

$$\hat{u}_\varepsilon^* \in [0, \hat{u}_{\varepsilon'}], \quad \hat{u}_\varepsilon^* \neq 0. \tag{3.9}$$

From (3.9), (3.6), (3.8) and the first part of the proof we infer that $\hat{u}_\varepsilon^* = \hat{u}_{\varepsilon'}$ and so, by (3.9), $\hat{u}_\varepsilon \leq \hat{u}_{\varepsilon'}$. This proves the Claim.

Next we will let $\varepsilon \rightarrow 0^+$ to produce a solution of the purely singular problem (3.1). To this end, let $\varepsilon_n \rightarrow 0^+$ and set $\hat{u}_n = \hat{u}_{\varepsilon_n}$ for all $n \in \mathbb{N}$. We have

$$\left\langle A_{p(\cdot)}(\hat{u}_n), h \right\rangle + \left\langle A_{q(\cdot)}(\hat{u}_n), h \right\rangle = \int_\Omega \frac{h}{[\hat{u}_n + \varepsilon_n]^{\eta(x)}} dx \tag{3.10}$$

for all $h \in W_0^{1,p(\cdot)}(\Omega)$ and for all $n \in \mathbb{N}$. Choosing $h = \hat{u}_n \in W_0^{1,p(\cdot)}(\Omega)$ leads to

$$\varrho_{p(\cdot)}(\nabla \hat{u}_n) \leq \int_\Omega \hat{u}_n^{1-\eta(x)} dx \quad \text{for all } n \in \mathbb{N}.$$

Therefore, $\{\hat{u}_n\}_{n \in \mathbb{N}} \subseteq W_0^{1,p(\cdot)}(\Omega)$ is bounded.

By passing to an appropriate subsequence if necessary, we may assume that

$$\hat{u}_n \xrightarrow{w} \bar{u} \quad \text{in } W_0^{1,p(\cdot)}(\Omega) \quad \text{and} \quad \hat{u}_n \rightarrow \bar{u} \quad \text{in } L^{p(\cdot)}(\Omega). \tag{3.11}$$

Now we choose $h = \hat{u}_n - \bar{u} \in W_0^{1,p(\cdot)}(\Omega)$. This yields

$$\begin{aligned} & \langle A_{p(\cdot)}(\hat{u}_n), \hat{u}_n - \bar{u} \rangle + \langle A_{q(\cdot)}(\hat{u}_n), \hat{u}_n - \bar{u} \rangle \\ &= \int_{\Omega} \frac{\hat{u}_n - \bar{u}}{[\hat{u}_n + \varepsilon_n]^{\eta(x)}} dx \leq \int_{\Omega} \frac{\hat{u}_n - \bar{u}}{\hat{u}_1^{\eta(x)}} dx \quad \text{for all } n \in \mathbb{N}, \end{aligned}$$

due to the Claim.

Let $\hat{d}(x) = \text{dist}(x, \partial\Omega)$ for all $x \in \bar{\Omega}$. Using Lemma 14.16 of Gilbarg-Trudinger [8, p. 355] we have that $\hat{d} \in \text{int}(C_0^1(\bar{\Omega})_+)$. We can find $c_3 > 0$ such that $c_3 \hat{d} \leq \hat{u}_1$, see Papageorgiou-Rădulescu-Repovš [11, p. 274]. Then we have for all $h \in W_0^{1,p(\cdot)}(\Omega)$ that

$$\left| \int_{\Omega} \frac{h}{\hat{u}_1^{\eta(x)}} dx \right| \leq c_4 \int_{\Omega} \frac{|h|}{\hat{d}} dx \leq c_5 \|\nabla h\|_{p(\cdot)}$$

for some $c_4, c_5 > 0$. Here we used the anisotropic Hardy inequality of Harjulehto-Hästö-Koskenoja [6]. From Marino-Winkert [10] (see also Ragusa-Tachikawa [19]) we have that $\{\hat{u}_n\}_{n \in \mathbb{N}} \subseteq L^\infty(\Omega)$ is bounded. Moreover by the lemma and its proof of Lazer-McKenna [9] we know that $\hat{u}_1^{-\eta(\cdot)} \in L^1(\Omega)$. So, from (3.11) and the dominated convergence theorem, it follows that

$$\int_{\Omega} \frac{\hat{u}_n - \bar{u}}{\hat{u}_1^{\eta(x)}} dx \longrightarrow 0 \quad \text{as } n \rightarrow \infty.$$

This implies

$$\limsup_{n \rightarrow \infty} [\langle A_{p(\cdot)}(\hat{u}_n), \hat{u}_n - \bar{u} \rangle + \langle A_{q(\cdot)}(\hat{u}_n), \hat{u}_n - \bar{u} \rangle] \leq 0,$$

which by the monotonicity of $A_{q(\cdot)}$ and the S_+ -property of $A_{p(\cdot)}$ (see Proposition 2.2 and the first part of the proof) leads to

$$\hat{u}_n \rightarrow \bar{u} \quad \text{in } W_0^{1,p(\cdot)}(\Omega) \quad \text{and} \quad \hat{u}_1 \leq \bar{u}. \tag{3.12}$$

So, if we pass to the limit in (3.10) as $n \rightarrow \infty$ and use the Lebesgue dominated convergence theorem, we then obtain

$$\langle A_{p(\cdot)}(\bar{u}), h \rangle + \langle A_{q(\cdot)}(\bar{u}), h \rangle = \int_{\Omega} \frac{h}{\bar{u}^{\eta(x)}} dx \quad \text{for all } h \in W_0^{1,p(\cdot)}(\Omega).$$

Since $\hat{u}_1 \leq \bar{u}$, we see that $\bar{u} \in W_0^{1,p(\cdot)}(\Omega)$ is a positive solution of (3.1). From Marino-Winkert [10] we know that $\bar{u} \in L^\infty(\Omega)$ and so we conclude that $\bar{u} \in \text{int}(C_0^1(\bar{\Omega})_+)$, see Zhang [25] and (3.12).

Finally, note that the function $\mathring{\mathbb{R}}_+ \ni s \rightarrow s^{-\eta(x)}$ is strictly decreasing. Therefore, the positive solution $\bar{u} \in \text{int}(C_0^1(\bar{\Omega})_+)$ is unique. □

In the next section we will use this solution to bypass the singularity and deal with C^1 -functionals on which we can apply the results of critical point theory.

4 Positive solutions

We introduce the following two sets

$$\begin{aligned} \mathcal{L} &= \{ \lambda > 0 : \text{problem } (P_\lambda) \text{ has a positive solution} \}, \\ \mathcal{S}_\lambda &= \{ u : u \text{ is a positive solution of problem } (P_\lambda) \}. \end{aligned}$$

Proposition 4.1 *If hypotheses H_0 and H_1 hold, then $\mathcal{L} \neq \emptyset$.*

Proof Let $\bar{u} \in \text{int}(C_0^1(\bar{\Omega})_+)$ be the unique positive solution of problem (3.1), see Proposition 3.1. By the anisotropic Hardy inequality, see Harjulehto-Hästö-Koskenoja [6], we know that $\bar{u}^{-\eta(\cdot)} h \in L^1(\Omega)$ for all $h \in W_0^{1,p(\cdot)}(\Omega)$. Hence, $\bar{u}^{-\eta(\cdot)} \in W^{1,p'(\cdot)}(\Omega) = W_0^{1,p(\cdot)}(\Omega)^*$.

We consider the following auxiliary Dirichlet problem

$$-\Delta_{p(\cdot)} u - \Delta_{q(\cdot)} u = \bar{u}^{-\eta(x)} + 1 \quad \text{in } \Omega, \quad u|_{\partial\Omega} = 0, \quad u > 0. \tag{4.1}$$

As in the proof of Proposition 3.1, exploiting the surjectivity and the strict monotonicity of the operator V , we infer that problem (4.1) admits a unique positive solution $\tilde{u} \in W_0^{1,p(\cdot)}(\Omega)$.

Since $\bar{u}^{-\eta(\cdot)} \leq c_6 \hat{d}^{-\eta(\cdot)}$ for some $c_6 > 0$, from Theorem B.1 of Saoudi-Ghanmi [20] we have

$$\tilde{u} \in \text{int}(C_0^1(\bar{\Omega})_+).$$

From the weak comparison principle, see Tolksdorf [24], we have that

$$\bar{u} \leq \tilde{u}. \tag{4.2}$$

Let $\lambda_0 = \frac{1}{\|N_f(\tilde{u})\|_\infty}$, see hypothesis H_1 (i). For $\lambda \in (0, \lambda_0]$ we have that

$$\lambda f(x, \tilde{u}(x)) \leq 1 \quad \text{for a. a. } x \in \Omega. \tag{4.3}$$

Applying (4.2) and (4.3) we get

$$-\Delta_{p(\cdot)} \tilde{u} - \Delta_{q(\cdot)} \tilde{u} = \bar{u}^{-\eta(x)} + 1 \geq \tilde{u}^{-\eta(x)} + \lambda f(x, \tilde{u}(x)) \quad \text{in } \Omega. \tag{4.4}$$

We introduce the Carathéodory function $i_\lambda : \Omega \times \overset{\circ}{\mathbb{R}}_+ \rightarrow \overset{\circ}{\mathbb{R}}_+$ defined by

$$i_\lambda(x, s) = \begin{cases} \bar{u}(x)^{-\eta(x)} + \lambda f(x, \bar{u}(x)) & \text{if } s < \bar{u}(x), \\ s^{-\eta(x)} + \lambda f(x, s) & \text{if } \bar{u}(x) \leq s \leq \tilde{u}(x), \\ \tilde{u}(x)^{-\eta(x)} + \lambda f(x, \tilde{u}(x)) & \text{if } \tilde{u}(x) < s. \end{cases} \tag{4.5}$$

We set $I_\lambda(x, s) = \int_0^s i_\lambda(x, t) dt$ and consider the C^1 -functional $\psi_\lambda : W_0^{1,p(\cdot)}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\psi_\lambda(u) = \int_\Omega \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_\Omega \frac{1}{q(x)} |\nabla u|^{q(x)} dx - \int_\Omega I_\lambda(x, u) dx$$

for all $u \in W_0^{1,p(\cdot)}(\Omega)$. Evidently, ψ_λ is coercive due to (4.5) and it is sequentially weakly lower semicontinuous. So, we can find $u_\lambda \in W_0^{1,p(\cdot)}(\Omega)$ such that

$$\psi_\lambda(u_\lambda) = \min \left[\psi_\lambda(u) : u \in W_0^{1,p(\cdot)}(\Omega) \right].$$

From this we know that $\psi'_\lambda(u_\lambda) = 0$ and so,

$$\langle A_{p(\cdot)}(u_\lambda), h \rangle + \langle A_{q(\cdot)}(u_\lambda), h \rangle = \int_\Omega i_\lambda(x, u_\lambda) h dx \tag{4.6}$$

for all $h \in W_0^{1,p(\cdot)}(\Omega)$. First we choose $h = (\bar{u} - u_\lambda)^+ \in W_0^{1,p(\cdot)}(\Omega)$ in (4.6). Then, by (4.5), $f \geq 0$ and Proposition 3.1 it follows that

$$\begin{aligned} & \langle A_{p(\cdot)}(u_\lambda), (\bar{u} - u_\lambda)^+ \rangle + \langle A_{q(\cdot)}(u_\lambda), (\bar{u} - u_\lambda)^+ \rangle \\ &= \int_\Omega i_\lambda(x, u_\lambda) (\bar{u} - u_\lambda)^+ dx \\ &= \int_\Omega \left[\bar{u}^{-\eta(x)} + \lambda f(x, \bar{u}) \right] (\bar{u} - u_\lambda)^+ dx \\ &\geq \int_\Omega \bar{u}^{-\eta(x)} (\bar{u} - u_\lambda)^+ dx \\ &= \langle A_{p(\cdot)}(\bar{u}), (\bar{u} - u_\lambda)^+ \rangle + \langle A_{q(\cdot)}(\bar{u}), (\bar{u} - u_\lambda)^+ \rangle. \end{aligned}$$

Therefore, $\bar{u} \leq u_\lambda$.

Next, we test (4.6) with $h = (u_\lambda - \tilde{u})^+ \in W_0^{1,p(\cdot)}(\Omega)$. As before, by (4.5) and (4.4), we have

$$\begin{aligned} & \langle A_{p(\cdot)}(u_\lambda), (u_\lambda - \tilde{u})^+ \rangle + \langle A_{q(\cdot)}(u_\lambda), (u_\lambda - \tilde{u})^+ \rangle \\ &= \int_\Omega i_\lambda(x, u_\lambda) (u_\lambda - \tilde{u})^+ dx \end{aligned}$$

$$\begin{aligned}
 &= \int_{\Omega} \left[\tilde{u}^{-\eta(x)} + \lambda f(x, \tilde{u}) \right] (u_{\lambda} - \tilde{u})^+ dx \\
 &\leq \langle A_{p(\cdot)}(\tilde{u}), (u_{\lambda} - \tilde{u})^+ \rangle + \langle A_{q(\cdot)}(\tilde{u}), (u_{\lambda} - \tilde{u})^+ \rangle.
 \end{aligned}$$

Hence, $u_{\lambda} \leq \tilde{u}$. So, we have proved that

$$u_{\lambda} \in [\bar{u}, \tilde{u}]. \tag{4.7}$$

Then, from (4.7), (4.5) and (4.6), it follows that

$$u_{\lambda} \in \mathcal{S}_{\lambda} \text{ for all } \lambda \in (0, \lambda_0].$$

Thus, $(0, \lambda_0] \subseteq \mathcal{L} \neq \emptyset$. □

We want to determine the regularity of the elements of the solution set \mathcal{S}_{λ} . To this end, we first establish a lower bound for the elements of \mathcal{S}_{λ} .

Proposition 4.2 *If hypotheses H_0, H_1 hold and $\lambda \in \mathcal{L}$, then $\bar{u} \leq u$ for all $u \in \mathcal{S}_{\lambda}$.*

Proof Let $u \in \mathcal{S}_{\lambda}$. We introduce the Carathéodory function $b: \Omega \times \overset{\circ}{\mathbb{R}}_+ \rightarrow \overset{\circ}{\mathbb{R}}_+$ defined by

$$b(x, s) = \begin{cases} s^{-\eta(x)} & \text{if } 0 < s < u(x), \\ u(x)^{-\eta(x)} & \text{if } u(x) < s. \end{cases} \tag{4.8}$$

We consider the following Dirichlet problem

$$-\Delta_{p(\cdot)} u - \Delta_{q(\cdot)} u = b(x, u) \text{ in } \Omega, \quad u|_{\partial\Omega} = 0, \quad u > 0. \tag{4.9}$$

As in the proof of Proposition 3.1, using approximations and fixed point theory, we can show that problem (4.9) has a positive solution $\bar{u}_0 \in W_0^{1,p(\cdot)}(\Omega)$. Applying (4.8), $f \geq 0$ and $u \in \mathcal{S}_{\lambda}$ yields

$$\begin{aligned}
 &\langle A_{p(\cdot)}(\bar{u}_0), (\bar{u}_0 - u)^+ \rangle + \langle A_{q(\cdot)}(\bar{u}_0), (\bar{u}_0 - u)^+ \rangle \\
 &= \int_{\Omega} b(x, \bar{u}_0) (\bar{u}_0 - u)^+ dx \\
 &= \int_{\Omega} u^{-\eta(x)} (\bar{u}_0 - u)^+ dx \\
 &\leq \int_{\Omega} \left[u^{-\eta(x)} + \lambda f(x, u) \right] (\bar{u}_0 - u)^+ dx \\
 &= \langle A_{p(\cdot)}(u), (\bar{u}_0 - u)^+ \rangle + \langle A_{q(\cdot)}(u), (\bar{u}_0 - u)^+ \rangle.
 \end{aligned}$$

Therefore, we have

$$\bar{u}_0 \leq u. \tag{4.10}$$

Then, (4.10), (4.8), (4.9) and Proposition 3.1 imply that

$$\bar{u}_0 = \bar{u} \in \text{int} \left(C_0^1(\bar{\Omega})_+ \right).$$

This shows that $\bar{u} \leq u$ for all $u \in \mathcal{S}_\lambda$, see (4.10). □

Using this lower bound and the anisotropic regularity theory of Saoudi-Ghanmi [20], we can have the regularity properties of the elements of \mathcal{S}_λ .

Proposition 4.3 *If hypotheses H_0, H_1 hold and $\lambda \in \mathcal{L}$, then $\emptyset \neq \mathcal{S}_\lambda \subseteq \text{int} \left(C_0^1(\bar{\Omega})_+ \right)$.*

Next we prove a structural property of \mathcal{L} , namely, we show that \mathcal{L} is connected, so an interval.

Proposition 4.4 *If hypotheses H_0, H_1 hold, $\lambda \in \mathcal{L}$ and $\mu \in (0, \lambda)$, then $\mu \in \mathcal{L}$.*

Proof Let $u_\lambda \in \mathcal{S}_\lambda \subseteq \text{int} \left(C_0^1(\bar{\Omega})_+ \right)$, see Proposition 4.3. We introduce the Carathéodory function $e_\mu : \Omega \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ defined by

$$e_\mu(x, s) = \begin{cases} \bar{u}(x)^{-\eta(x)} + \mu f(x, \bar{u}(x)) & \text{if } s < \bar{u}(x), \\ s^{-\eta(x)} + \mu f(x, s) & \text{if } \bar{u}(x) \leq s \leq u_\lambda(x), \\ u_\lambda(x)^{-\eta(x)} + \mu f(x, u_\lambda(x)) & \text{if } u_\lambda(x) < s. \end{cases} \quad (4.11)$$

We set $E_\mu(x, s) = \int_0^s e_\mu(x, t) dt$ and consider the C^1 -functional $\sigma_\mu : W_0^{1,p(\cdot)}(\Omega) \rightarrow \mathbb{R}$ defined by

$$\sigma_\mu(u) = \int_\Omega \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_\Omega \frac{1}{q(x)} |\nabla u|^{q(x)} dx - \int_\Omega E_\mu(x, u) dx$$

for all $u \in W_0^{1,p(\cdot)}(\Omega)$. It is clear that σ_μ is coercive because of (4.11) and it is sequentially weakly lower semicontinuous. So, there exists $u_\mu \in W_0^{1,p(\cdot)}(\Omega)$ such that

$$\sigma_\mu(u_\mu) = \min \left[\sigma_\mu(u) : u \in W_0^{1,p(\cdot)}(\Omega) \right].$$

That means $\sigma'_\mu(u_\mu) = 0$ and so,

$$\langle A_{p(\cdot)}(u_\mu), h \rangle + \langle A_{q(\cdot)}(u_\mu), h \rangle = \int_\Omega e_\mu(x, u_\mu) h dx \quad (4.12)$$

for all $h \in W_0^{1,p(\cdot)}(\Omega)$. If we choose $h = (\bar{u} - u_\mu)^+ \in W_0^{1,p(\cdot)}(\Omega)$ in (4.12) we can show that $\bar{u} \leq u_\mu$, see the proof of Proposition 4.1. Next, we choose $h = (u_\mu - u_\lambda)^+ \in W_0^{1,p(\cdot)}(\Omega)$ in (4.12). Then, by (4.11), $f \geq 0, \mu < \lambda$ and $u_\lambda \in \mathcal{S}_\lambda$, we obtain

$$\begin{aligned} & \left\langle A_{p(\cdot)}(u_\mu), (u_\mu - u_\lambda)^+ \right\rangle + \left\langle A_{q(\cdot)}(u_\mu), (u_\mu - u_\lambda)^+ \right\rangle \\ &= \int_\Omega e_\mu(x, u_\mu) (u_\mu - u_\lambda)^+ dx \\ &= \int_\Omega \left[u_\lambda^{-\eta(x)} + \mu f(x, u_\lambda) \right] (u_\mu - u_\lambda)^+ dx \\ &\leq \int_\Omega \left[u_\lambda^{-\eta(x)} + \lambda f(x, u_\lambda) \right] (u_\mu - u_\lambda)^+ dx \\ &= \left\langle A_{p(\cdot)}(u_\lambda), (u_\mu - u_\lambda)^+ \right\rangle + \left\langle A_{q(\cdot)}(u_\lambda), (u_\mu - u_\lambda)^+ \right\rangle. \end{aligned}$$

Hence, $u_\mu \leq u_\lambda$. Therefore we have

$$u_\mu \in [\bar{u}, u_\lambda]. \tag{4.13}$$

From (4.13), (4.11) and (4.12) it follows that

$$u_\mu \in \mathcal{S}_\mu \subseteq \text{int} \left(C_0^1(\bar{\Omega})_+ \right) \quad \text{and so} \quad \mu \in \mathcal{L}.$$

□

From Proposition 4.4 and its proof we have the following corollary.

Corollary 4.5 *If hypotheses H_0, H_1 hold and if $\lambda \in \mathcal{L}, u_\lambda \in \mathcal{S}_\lambda \subseteq \text{int} (C_0^1(\bar{\Omega})_+)$ and $0 < \mu < \lambda$, then $\mu \in \mathcal{L}$ and there exists $u_\mu \in \mathcal{S}_\mu \subseteq \text{int} (C_0^1(\bar{\Omega})_+)$ such that $u_\mu \leq u_\lambda$.*

In the next proposition we are going to improve the assertion of Corollary 4.5.

Proposition 4.6 *If hypotheses H_0, H_1 hold and if $\lambda \in \mathcal{L}, u_\lambda \in \mathcal{S}_\lambda \subseteq \text{int} (C_0^1(\bar{\Omega})_+)$ and $0 < \mu < \lambda$, then $\mu \in \mathcal{L}$ and there exists $u_\mu \in \mathcal{S}_\mu \subseteq \text{int} (C_0^1(\bar{\Omega})_+)$ such that*

$$u_\lambda - u_\mu \in \text{int} \left(C_0^1(\bar{\Omega})_+ \right).$$

Proof From Corollary 4.5 we already know that $\mu \in \mathcal{L}$ and that there exists $u_\mu \in \mathcal{S}_\mu \subseteq \text{int} (C_0^1(\bar{\Omega})_+)$ such that

$$u_\mu \leq u_\lambda. \tag{4.14}$$

Let $\rho = \|u_\lambda\|_\infty$ and let $\hat{\xi}_\rho > 0$ be as postulated by hypothesis H_1 (iv). Applying $u_\mu \in \mathcal{S}_\mu$, (4.14), hypothesis H_1 (iv), $f \geq 0, \mu < \lambda$ and $u_\lambda \in \mathcal{S}_\lambda$ gives

$$\begin{aligned} & -\Delta_{p(\cdot)}u_\mu - \Delta_{q(\cdot)}u_\mu + \mu \hat{\xi}_\rho u_\mu^{p(x)-1} - u_\mu^{-\eta(x)} \\ &= \mu \left[f(x, u_\mu) + \hat{\xi}_\rho u_\mu^{p(x)-1} \right] \\ &\leq \mu \left[f(x, u_\lambda) + \hat{\xi}_\rho u_\lambda^{p(x)-1} \right] \end{aligned}$$

$$\begin{aligned} &\leq \lambda f(x, u_\lambda) + \mu \hat{\xi}_\rho u_\lambda^{p(x)-1} \\ &= -\Delta_{p(\cdot)} u_\lambda - \Delta_{q(\cdot)} u_\lambda + \mu \hat{\xi}_\rho u_\lambda^{p(x)-1} - u_\lambda^{-\eta(x)}. \end{aligned} \tag{4.15}$$

Note that since $u_\mu \in \text{int}(C_0^1(\overline{\Omega})_+)$, $f \geq 0$ and $\mu < \lambda$, we have

$$(\lambda - \mu) \left[N_f(u_\mu) + \hat{\xi}_\rho u_\mu^{p(\cdot)-1} \right] \succeq 0.$$

Hence, from (4.15) and Proposition 2.3(a), we infer that

$$u_\lambda - u_\mu \in \text{int}(C_0^1(\overline{\Omega})_+).$$

□

We set $\lambda^* = \sup \mathcal{L}$.

Proposition 4.7 *If hypotheses H_0, H_1 hold, then $\lambda^* < +\infty$.*

Proof Hypotheses H_1 (i), (ii) and (iii) imply that we can find $\hat{\lambda} > 0$ such that

$$s^{-\eta(x)} + \hat{\lambda} f(x, s) \geq s^{p(x)-1} \quad \text{for a. a. } x \in \Omega \text{ and for all } s > 0. \tag{4.16}$$

Let $\lambda > \hat{\lambda}$ and suppose that $\lambda \in \mathcal{L}$. We can find $u \in \mathcal{S}_\lambda \subseteq \text{int}(C_0^1(\overline{\Omega})_+)$ and from Proposition 4.2 we have $\bar{u} \leq u$. Let $\Omega_0 \subseteq \Omega$ be an open subset with C^2 -boundary, $\overline{\Omega}_0 \subseteq \Omega$ and $m_0 = \min_{x \in \overline{\Omega}_0} u(x) \leq 1$. Note that since $u \in \text{int}(C_0^1(\overline{\Omega})_+)$ we have $0 < m_0$. Let $\delta \in (0, 1)$ be small and set $m_0^\delta = m_0 + \delta$. Note that

$$0 \leq \frac{1}{m_0^{\eta(x)}} - \frac{1}{(m_0^\delta)^{\eta(x)}} \leq \frac{\delta \eta(x)}{m_0^{2\eta(x)}} \leq \frac{\delta \eta^-}{m_0^{2\eta_+}} \quad \text{for all } x \in \overline{\Omega}. \tag{4.17}$$

Let $\rho = \|u\|_\infty$ and let $\hat{\xi}_\rho > 0$ be as postulated by hypothesis H_1 (iv). Then, by applying (4.17), (4.16), $m_0 \leq 1$, $\delta > 0$ small enough, $f \geq 0$ and $\lambda > \hat{\lambda}$ we obtain

$$\begin{aligned} &-\Delta_{p(\cdot)} m_0^\delta - \Delta_{q(\cdot)} m_0^\delta + \hat{\lambda} \hat{\xi}_\rho (m_0^\delta)^{p(x)-1} - (m_0^\delta)^{-\eta(x)} \\ &\leq \hat{\lambda} \hat{\xi}_\rho m_0^{p(x)-1} + \chi(\delta) - m_0^{-\eta(x)} \quad \text{with } \chi(\delta) \rightarrow 0^+ \text{ as } \delta \rightarrow 0^+, \\ &\leq \left[\hat{\lambda} \hat{\xi}_\rho + 1 \right] m_0^{p(x)-1} + \chi(\delta) - m_0^{-\eta(x)} \\ &\leq \hat{\lambda} \left[f(x, m_0) + \hat{\xi}_\rho m_0^{p(x)-1} \right] + \chi(\delta) - m_0^{-\eta_+} \\ &< \hat{\lambda} \left[f(x, u) + \hat{\xi}_\rho u^{p(x)-1} \right] \\ &\leq \lambda f(x, u) + \hat{\lambda} \hat{\xi}_\rho u^{p(x)-1} \\ &= -\Delta_{p(\cdot)} u - \Delta_{q(\cdot)} u + \hat{\lambda} \hat{\xi}_\rho u^{p(x)-1} - u^{-\eta(x)} \quad \text{in } \Omega_0. \end{aligned}$$

Then, by Proposition 2.3(b), we get $m_0^\delta < u(x)$ for all $x \in \Omega_0$ and for all $\delta \in (0, 1)$ small enough. This contradicts the definition of m_0 . Therefore, $\lambda^* \leq \hat{\lambda} < +\infty$. \square

Next we are going to prove that we have multiple solutions for all $\lambda \in (0, \lambda^*)$.

Proposition 4.8 *If hypotheses H_0, H_1 hold and $\lambda \in (0, \lambda^*)$, then problem (P_λ) has at least two positive solutions*

$$u_0, \hat{u} \in \text{int} \left(C_0^1(\bar{\Omega})_+ \right) \text{ with } u_0 \neq \hat{u}.$$

Proof Let $0 < \lambda < \vartheta < \lambda^*$. On account of Proposition 4.6 we can find $u_\vartheta \in \mathcal{S}_\vartheta \subseteq \text{int} \left(C_0^1(\bar{\Omega})_+ \right)$ and $u_0 \in \mathcal{S}_\lambda \subseteq \text{int} \left(C_0^1(\bar{\Omega})_+ \right)$ such that

$$u_\vartheta - u_0 \in \text{int} \left(C_0^1(\bar{\Omega})_+ \right). \tag{4.18}$$

Also from Proposition 4.2 we have

$$\bar{u} \leq u_0. \tag{4.19}$$

Let $\rho = \|u_0\|_\infty$ and let $\hat{\xi}_\rho > 0$ be as postulated by hypothesis H_1 (iv). Then, using $f \geq 0$, (4.19), hypothesis H_1 (iv) and $u_0 \in \mathcal{S}_\lambda$, we obtain

$$\begin{aligned} & -\Delta_{p(\cdot)}\bar{u} - \Delta_{q(\cdot)}\bar{u} + \lambda\hat{\xi}_\rho\bar{u}^{p(x)-1} - \bar{u}^{-\eta(x)} \\ & \leq \lambda \left[f(x, \bar{u}) + \hat{\xi}_\rho\bar{u}^{p(x)-1} \right] \\ & \leq \lambda \left[f(x, u_0) + \hat{\xi}_\rho u_0^{p(x)-1} \right] \\ & = -\Delta_{p(\cdot)}u_0 - \Delta_{q(\cdot)}u_0 + \lambda\hat{\xi}_\rho u_0^{p(x)-1} - u_0^{-\eta(x)} \quad \text{in } \Omega. \end{aligned} \tag{4.20}$$

Note that $0 \leq \hat{\xi}_\rho u_0^{p(x)-1}$ since $u_0 \in \text{int} \left(C_0^1(\bar{\Omega})_+ \right)$. So, from (4.20) and Proposition 2.3(a), we get that

$$u_0 - \bar{u} \in \text{int} \left(C_0^1(\bar{\Omega})_+ \right). \tag{4.21}$$

From (4.18) and (4.21) it follows that

$$u_0 \in \text{int}_{C_0^1(\bar{\Omega})} [\bar{u}, u_\vartheta]. \tag{4.22}$$

We introduce the Carathéodory function $j_\lambda : \Omega \times \overset{\circ}{\mathbb{R}}_+ \rightarrow \overset{\circ}{\mathbb{R}}_+$ defined by

$$j_\lambda(x, s) = \begin{cases} \bar{u}(x)^{-\eta(x)} + \lambda f(x, \bar{u}(x)) & \text{if } s \leq \bar{u}(x), \\ s^{-\eta(x)} + \lambda f(x, s) & \text{if } \bar{u}(x) < s. \end{cases} \tag{4.23}$$

Moreover, we introduce the truncation of $j_\lambda(x, \cdot)$ at $u_\vartheta(x)$, namely, the Carathéodory function $\hat{j}_\lambda : \Omega \times \overset{\circ}{\mathbb{R}}_+ \rightarrow \overset{\circ}{\mathbb{R}}_+$ defined by

$$\hat{j}_\lambda(x, s) = \begin{cases} j_\lambda(x, s) & \text{if } s \leq u_\vartheta(x), \\ j_\lambda(x, u_\vartheta(x)) & \text{if } u_\vartheta(x) < s. \end{cases} \tag{4.24}$$

We set $J_\lambda(x, s) = \int_0^s j_\lambda(x, t) dt$ and $\hat{J}_\lambda(x, s) = \int_0^s \hat{j}_\lambda(x, t) dt$ and consider the C^1 -functionals $w_\lambda, \hat{w}_\lambda : W_0^{1,p(\cdot)}(\Omega) \rightarrow \mathbb{R}$ defined by

$$w_\lambda(u) = \int_\Omega \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_\Omega \frac{1}{q(x)} |\nabla u|^{q(x)} dx - \int_\Omega J_\lambda(x, u) dx,$$

$$\hat{w}_\lambda(u) = \int_\Omega \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_\Omega \frac{1}{q(x)} |\nabla u|^{q(x)} dx - \int_\Omega \hat{J}_\lambda(x, u) dx$$

for all $u \in W_0^{1,p(\cdot)}(\Omega)$.

From (4.23) and (4.24) it is clear that

$$w_\lambda|_{[0, u_\vartheta]} = \hat{w}_\lambda|_{[0, u_\vartheta]} \quad \text{and} \quad w'_\lambda|_{[0, u_\vartheta]} = \hat{w}'_\lambda|_{[0, u_\vartheta]}. \tag{4.25}$$

Moreover, applying (4.23) and (4.24), we can easily show that

$$K_{w_\lambda} \subseteq [\bar{u}] \cap \text{int} \left(C_0^1(\bar{\Omega})_+ \right) \quad \text{and} \quad K_{\hat{w}_\lambda} \subseteq [\bar{u}, u_\vartheta] \cap \text{int} \left(C_0^1(\bar{\Omega})_+ \right). \tag{4.26}$$

On account of (4.24) and (4.26), we see that we may assume that

$$K_{\hat{w}_\lambda} = \{u_0\}. \tag{4.27}$$

Otherwise we already have a second positive smooth solution for problem (P_λ) and so we are done, see (4.24) and (4.26).

From (4.24) we see that the functional $\hat{w}_\lambda : W_0^{1,p(\cdot)}(\Omega) \rightarrow \mathbb{R}$ is coercive and it is easy to check that it is sequentially weakly lower semicontinuous. Hence, its global minimizer $\hat{u}_0 \in W_0^{1,p(\cdot)}(\Omega)$ exists, that is,

$$\hat{w}_\lambda(\hat{u}_0) = \min \left[\hat{w}_\lambda(u) : u \in W_0^{1,p(\cdot)}(\Omega) \right].$$

From (4.27) we conclude that $\hat{u}_0 = u_0$. From (4.22) and (4.25) it follows that u_0 is a local $C_0^1(\bar{\Omega})$ -minimizer of w_λ , Hence

$$u_0 \text{ is a local } W_0^{1,p(\cdot)}(\Omega) \text{-minimizer of } w_\lambda, \tag{4.28}$$

see Tan-Fang [23] and Gasiński-Papageorgiou [7]. From (4.23) and (4.26) we see that we can assume that

$$K_{w_\lambda} \text{ is finite.} \tag{4.29}$$

Otherwise we already have an infinity of positive smooth solutions for problem (P_λ) and so we are done.

Then, from (4.28), (4.29) and Theorem 5.7.4 of Papageorgiou-Rădulescu-Repovš [12, p. 449] we know that there exists $\rho \in (0, 1)$ small such that

$$w_\lambda(u_0) < \inf [w_\lambda(u) : \|u - u_0\| = \rho] = m_\lambda. \tag{4.30}$$

On account of hypothesis $H_1(ii)$, if $u \in \text{int}(C_0^1(\bar{\Omega})_+)$, then

$$w_\lambda(tu) \rightarrow -\infty \text{ as } t \rightarrow +\infty. \tag{4.31}$$

In order to apply the mountain pass theorem we only need to show that the functional w_λ satisfies the C-condition.

Claim: w_λ fulfills the C-condition.

We consider the sequence $\{u_n\}_{n \in \mathbb{N}} \subseteq W_0^{1,p(\cdot)}(\Omega)$ such that

$$|w_\lambda(u_n)| \leq c_7 \text{ for some } c_7 > 0 \text{ and for all } n \in \mathbb{N}, \tag{4.32}$$

$$(1 + \|u_n\|)w'_\lambda(u_n) \rightarrow 0 \text{ in } W^{-1,p'(\cdot)}(\Omega) \text{ as } n \rightarrow \infty. \tag{4.33}$$

From (4.33) we have

$$\left| \langle A_{p(\cdot)}(u_n), h \rangle + \langle A_{q(\cdot)}(u_n), h \rangle - \int_\Omega j_\lambda(x, u_n)h \, dx \right| \leq \frac{\varepsilon_n \|h\|}{1 + \|u_n\|} \tag{4.34}$$

for all $h \in W_0^{1,p(\cdot)}(\Omega)$ with $\varepsilon_n \rightarrow 0^+$. Choosing $h = -u_n^- \in W_0^{1,p(\cdot)}(\Omega)$ in (4.34), recalling that $\bar{u}^{-\eta(\cdot)}h \in L^1(\Omega)$ for all $h \in W_0^{1,p(\cdot)}(\Omega)$, see Harjulehto-Hästö-Koskenoja [6], and applying (4.23) leads to

$$\varrho_{p(\cdot)}(\nabla u_n^-) + \varrho_{q(\cdot)}(\nabla u_n^-) \leq c_8 \|u_n^-\| \text{ for some } c_8 > 0 \text{ and for all } n \in \mathbb{N},$$

which implies that

$$\{u_n^-\}_{n \in \mathbb{N}} \subseteq W_0^{1,p(\cdot)}(\Omega) \text{ is bounded.} \tag{4.35}$$

Now we choose $h = u_n^+ \in W_0^{1,p(\cdot)}(\Omega)$ as test function in (4.34). This gives

$$-\varrho_{p(\cdot)}(\nabla u_n^+) - \varrho_{q(\cdot)}(\nabla u_n^+) + \int_\Omega j_\lambda(x, u_n^+) u_n^+ \, dx \leq \varepsilon_n \text{ for all } n \in \mathbb{N}. \tag{4.36}$$

Furthermore, from (4.32) and (4.35), we obtain

$$\left| \int_\Omega \frac{1}{p(x)} |\nabla u_n^+|^{p(x)} \, dx + \int_\Omega \frac{1}{q(x)} |\nabla u_n^+|^{q(x)} \, dx - \int_\Omega J_\lambda(x, u_n^+) \, dx \right| \leq c_9$$

for some $c_9 > 0$ and for all $n \in \mathbb{N}$. This implies

$$\varrho_{p(\cdot)}(\nabla u_n^+) + \varrho_{q(\cdot)}(\nabla u_n^+) - \int_{\Omega} p_+ J_{\lambda}(x, u_n^+) dx \leq p_+ c_9 \quad \text{for all } n \in \mathbb{N}. \quad (4.37)$$

We add (4.36) and (4.37) and obtain

$$\int_{\Omega} [j_{\lambda}(x, u_n^+) u_n^+ - p_+ J_{\lambda}(x, u_n^+)] dx \leq c_{10} \quad \text{for some } c_{10} > 0 \text{ and for all } n \in \mathbb{N},$$

which by (4.23) results in

$$\int_{\Omega} \lambda [f(x, u_n^+) u_n^+ - p_+ F(x, u_n^+)] dx \leq c_{11} \left(1 + \int_{\Omega} (u_n^+)^{1-\eta(x)} dx \right) \quad (4.38)$$

for some $c_{11} > 0$ and for all $n \in \mathbb{N}$.

Hypotheses $H_1(i)$, (iii) imply the existence of $\gamma_1 \in (0, \gamma_0)$ and $c_{12} > 0$ such that

$$\gamma_1 s^{-\tau(x)} - c_{12} \leq f(x, s)s - p_+ F(x, s) \quad \text{for a. a. } x \in \Omega \text{ and for all } s \geq 0. \quad (4.39)$$

Using (4.39) in (4.38), we have

$$\varrho_{\tau(\cdot)}(u_n^+) \leq c_{13} \left[1 + \int_{\Omega} (u_n^+)^{1-\eta(x)} dx \right] \quad \text{for some } c_{13} > 0 \text{ and for all } n \in \mathbb{N}.$$

Hence, we see that

$$\{u_n^+\}_{n \in \mathbb{N}} \subseteq L^{\tau(\cdot)}(\Omega) \text{ is bounded.} \quad (4.40)$$

From hypothesis $H_1(iii)$ we see that, without any loss of generality, we may assume that $\tau(x) < r < p_-^*$ for all $x \in \overline{\Omega}$. Hence, $\tau_- < r < p_-^*$ and so we can find $t \in (0, 1)$ such that

$$\frac{1}{r} = \frac{1-t}{\tau_-} + \frac{t}{p_-^*}. \quad (4.41)$$

Applying the interpolation inequality, see Papageorgiou-Winkert [15, p. 116], we have

$$\|u_n^+\|_r \leq \|u_n^+\|_{\tau_-}^{1-t} \|u_n^+\|_{p_-^*}^t.$$

Thus, due to (4.40),

$$\|u_n^+\|_r^t \leq c_{14} \|u_n^+\|_{p_-^*}^{tr} \quad \text{for some } c_{14} > 0 \text{ and for all } n \in \mathbb{N}.$$

Then, by the Sobolev embedding theorem, we obtain

$$\|u_n^+\|_r^r \leq c_{15} \|u_n^+\|^{tr} \quad \text{for some } c_{15} > 0 \text{ and for all } n \in \mathbb{N}. \tag{4.42}$$

We take $h = u_n^+ \in W_0^{1,p(\cdot)}(\Omega)$ in (4.34) as test function and get

$$\varrho_{p(\cdot)}(\nabla u_n^+) + \varrho_{q(\cdot)}(\nabla u_n^+) \leq \varepsilon_n + \int_{\Omega} j_{\lambda}(x, u_n^+) u_n^+ dx \quad \text{for all } n \in \mathbb{N},$$

which by (4.23) and (4.42) gives

$$\begin{aligned} \varrho_{p(\cdot)}(\nabla u_n^+) + \varrho_{q(\cdot)}(\nabla u_n^+) &\leq c_{16} \left[1 + \int_{\Omega} \lambda f(x, u_n^+) u_n^+ dx \right] \\ &\leq c_{17} [1 + \lambda \|u_n^+\|_r^r] \\ &\leq c_{18} [1 + \lambda \|u_n^+\|^{tr}] \end{aligned} \tag{4.43}$$

for some $c_{16}, c_{17}, c_{18} > 0$ and for all $n \in \mathbb{N}$.

From (4.41) we have

$$tr = \frac{p_-^* (r - \tau_-)}{p_-^* - \tau_-} < p_-.$$

Therefore, from (4.43) and Proposition 2.1 it follows that

$$\{u_n\}_{n \in \mathbb{N}} \subseteq W_0^{1,p(\cdot)}(\Omega) \text{ is bounded.}$$

So, we may assume that

$$u_n \xrightarrow{w} u \text{ in } W_0^{1,p(\cdot)}(\Omega) \text{ and } u_n \rightarrow u \text{ in } L^{p(\cdot)}(\Omega). \tag{4.44}$$

We choose $h = u_n - u \in W_0^{1,p(\cdot)}(\Omega)$ in (4.34), pass to the limit as $n \rightarrow \infty$ and apply (4.44). This yields

$$\lim_{n \rightarrow \infty} [\langle A_{p(\cdot)}(u_n), u_n - u \rangle + \langle A_{q(\cdot)}(u_n), u_n - u \rangle] = 0.$$

Note that $A_{q(\cdot)}(\cdot)$ is monotone, so we have

$$\limsup_{n \rightarrow \infty} [\langle A_{p(\cdot)}(u_n), u_n - u \rangle + \langle A_{q(\cdot)}(u), u_n - u \rangle] \leq 0.$$

Because of (4.44) we then derive

$$\limsup_{n \rightarrow \infty} \langle A_{p(\cdot)}(u_n), u_n - u \rangle \leq 0$$

and so, by Proposition 2.2,

$$u_n \rightarrow u \text{ in } W_0^{1,p(\cdot)}(\Omega).$$

This proves the Claim.

Then, (4.30), (4.31) and the Claim permit us the use of the mountain pass theorem. So we can find $\hat{u} \in W_0^{1,p(\cdot)}(\Omega)$ such that

$$\hat{u} \in K_{w_\lambda} \subseteq [\bar{u}] \cap \text{int} \left(C_0^1(\bar{\Omega})_+ \right),$$

see (4.26), and

$$w_\lambda(u_0) < m_\lambda \leq w_\lambda(\hat{u}),$$

see (4.30). We conclude that $\hat{u} \in \text{int} \left(C_0^1(\bar{\Omega})_+ \right)$ is the second positive solution of (P_λ) for $\lambda \in (0, \lambda^*)$ and $\hat{u} \neq u_0$. \square

It remains to decide whether the critical parameter value $\lambda^* > 0$ is admissible.

Proposition 4.9 *If hypotheses H_0 and H_1 hold, then $\lambda^* \in \mathcal{L}$.*

Proof Let $\{\lambda_n\}_{n \in \mathbb{N}} \subseteq (0, \lambda^*) \subseteq \mathcal{L}$ be such that $\lambda_n \nearrow \lambda^*$ as $n \rightarrow \infty$. From the proof of Proposition 3.10 we know that we can find $u_n \in \mathcal{S}_{\lambda_n} \subseteq \text{int} \left(C_0^1(\bar{\Omega})_+ \right)$ such that

$$w_{\lambda_n}(u_n) \leq w_{\lambda_n}(\bar{u}) \text{ for all } n \in \mathbb{N}.$$

Applying (4.23), $f \geq 0$ and Proposition 3.1 we obtain

$$\begin{aligned} & w_{\lambda_n}(u_n) \\ & \leq \frac{1}{q_-} \left[\varrho_{p(\cdot)}(\nabla \bar{u}) + \varrho_{q(\cdot)}(\nabla \bar{u}) - \int_{\Omega} \bar{u}^{1-\eta(x)} dx - \int_{\Omega} \lambda_n f(x, \bar{u}) \bar{u} dx \right] \\ & \leq \frac{1}{q_-} \left[\varrho_{p(\cdot)}(\nabla \bar{u}) + \varrho_{q(\cdot)}(\nabla \bar{u}) \right] - \int_{\Omega} \bar{u}^{1-\eta(x)} dx \\ & \leq \left[\frac{1}{q_-} - 1 \right] \left(\varrho_{p(\cdot)}(\nabla \bar{u}) + \varrho_{q(\cdot)}(\nabla \bar{u}) \right) < 0 \end{aligned} \tag{4.45}$$

for all $n \in \mathbb{N}$. Furthermore, we have

$$\langle A_{p(\cdot)}(u_n), h \rangle + \langle A_{q(\cdot)}(u_n), h \rangle = \int_{\Omega} j_\lambda(x, u_n) h dx \tag{4.46}$$

for all $h \in W_0^{1,p(\cdot)}(\Omega)$ and for all $n \in \mathbb{N}$.

Using (4.45) and (4.46) and reasoning as in the Claim in the proof of Proposition 4.8, we obtain

$$u_n \rightarrow u^* \text{ in } W_0^{1,p(\cdot)}(\Omega) \text{ and } \bar{u} \leq u^*,$$

see Proposition 4.2. Hence, $u^* \in \mathcal{S}_{\lambda^*} \subseteq \text{int}(C_0^1(\bar{\Omega})_+)$ and so $\lambda^* \in \mathcal{L}$. □

So, we have proved that

$$\mathcal{L} = (0, \lambda^*].$$

Summarizing our results we can state the following bifurcation-type result describing the changes in the set of positive solutions as the parameter moves on $\mathring{\mathbb{R}}_+ = (0, +\infty)$.

Theorem 4.10 *If hypotheses H_0 and H_1 hold, then there exists $\lambda^* > 0$ such that*

(a) *for every $\lambda \in (0, \lambda^*)$, problem (P_λ) has at least two positive solutions*

$$u_0, \hat{u} \in \text{int}(C_0^1(\bar{\Omega})_+), \quad u_0 \neq \hat{u};$$

(b) *for $\lambda = \lambda^*$, problem (P_λ) has at least one positive solution*

$$u^* \in \text{int}(C_0^1(\bar{\Omega})_+);$$

(c) *for every $\lambda > \lambda^*$, problem (P_λ) has no positive solutions.*

Acknowledgements The authors wish to thank the two anonymous referees for their constructive remarks.

Funding Open Access funding enabled and organized by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Bahrouni, A., Radulescu, V.D., Repovš, D.: Double phase transonic flow problems with variable growth: nonlinear patterns and stationary waves. *Nonlinearity* **32**(7), 2481–2495 (2019)
2. Byun, S. S., Ko, E.: Global $C^{1,\alpha}$ regularity and existence of multiple solutions for singular $p(x)$ -Laplacian equations. *Calc. Var. Partial Differential Equations* **56** (2017), no. 5, Paper No. 76, 29 pp
3. Cherfils, L., Il'yasov, Y.: On the stationary solutions of generalized reaction diffusion equations with p - q -Laplacian. *Commun. Pure Appl. Anal.* **4**(1), 9–22 (2005)

4. Diening, L., Harjulehto, P., Hasto, P., Ruzicka, M.: Lebesgue and Sobolev spaces with variable exponents. Springer, Heidelberg (2011)
5. Fan, X., Zhao, D.: A class of De Giorgi type and Hölder continuity. *Nonlinear Anal.* **36**(3), 295–318 (1999)
6. Harjulehto, P., Hästö, P., Koskenoja, M.: Hardy's inequality in a variable exponent Sobolev space. *Georgian Math. J.* **12**(3), 431–442 (2005)
7. Gasiński, L., Papageorgiou, N.S.: Anisotropic nonlinear Neumann problems. *Calc. Var. Partial Differ. Equ.* **42**(3–4), 323–354 (2011)
8. Gilbarg, D., Trudinger, N.S.: *Elliptic Partial Differential Equations of Second Order*. Springer, Berlin (2001)
9. Lazer, A.C., McKenna, P.J.: On a singular nonlinear elliptic boundary-value problem. *Proc. Amer. Math. Soc.* **111**(3), 721–730 (1991)
10. Marino, G., Winkert, P.: L^∞ -bounds for general singular elliptic equations with convection term. *Appl. Math. Lett.* **107**, 106410, 6 pp (2020)
11. Papageorgiou, N.S., Rădulescu, V.D., Repovš, D.D.: Anisotropic equations with indefinite potential and competing nonlinearities. *Nonlinear Anal.* **201**, 111861 (2020)
12. Papageorgiou, N.S., Rădulescu, V.D., Repovš, D.D.: *Nonlinear Analysis – Theory and Methods*. Springer, Cham (2019)
13. Papageorgiou, N.S., Rădulescu, V.D., Repovš, D.D.: Nonlinear nonhomogeneous singular problems, *Calc. Var. Partial Differential Equations* **59** (2020), no. 1, Paper No. 9, 31 pp
14. Papageorgiou, N.S., Vetro, C., Vetro, F.: Multiple solutions for parametric double phase Dirichlet problems. *Commun. Contemp. Math.* **23**(4), 2050006 (2021)
15. Papageorgiou, N.S., Winkert, P.: *Applied Nonlinear Functional Analysis. an Introduction*. De Gruyter, Berlin (2018)
16. Papageorgiou, N.S., Winkert, P.: Singular p -Laplacian equations with superlinear perturbation. *J. Diff. Equ.* **266**(2–3), 1462–1487 (2019)
17. Papageorgiou, N.S., Zhang, C.: Noncoercive resonant $(p, 2)$ -equations with concave terms. *Adv. Nonlinear Anal.* **9**(1), 228–249 (2020)
18. Rădulescu, V.D., Repovš, D.D.: *Partial Differential Equations with Variable Exponents*. CRC Press, Boca Raton (2015)
19. Ragusa, M.A., Tachikawa, A.: Regularity for minimizers for functionals of double phase with variable exponents. *Adv. Nonlinear Anal.* **9**(1), 710–728 (2020)
20. Saoudi, K., Ghanmi, A.: A multiplicity results for a singular equation involving the $p(x)$ -Laplace operator. *Complex Var. Elliptic Equ.* **62**(5), 695–725 (2017)
21. Saoudi, K., Kratou, M., Alsadhan, S.: Multiplicity results for the $p(x)$ -Laplacian equation with singular nonlinearities and nonlinear Neumann boundary condition, *Int. J. Differ. Equ.* **2016**, Art. ID 3149482, 14 pp
22. Struwe, M.: *Variational Methods*. Springer-Verlag, Berlin, fourth edition (2008)
23. Tan, Z., Fang, F.: Orlicz-Sobolev versus Hölder local minimizer and multiplicity results for quasilinear elliptic equations. *J. Math. Anal. Appl.* **402**(1), 348–370 (2013)
24. Tolksdorf, P.: On the Dirichlet problem for quasilinear equations in domains with conical boundary points. *Comm. Partial Differ. Equ.* **8**(7), 773–817 (1983)
25. Zhang, Q.: A strong maximum principle for differential equations with nonstandard $p(x)$ -growth conditions. *J. Math. Anal. Appl.* **312**(1), 24–32 (2005)
26. Zhikov, V.V.: On variational problems and nonlinear elliptic equations with nonstandard growth conditions. *J. Math. Sci. (N. Y.)* **173**(5), 463–570 (2011)